

# A CLASS OF FOURIER MULTIPLIERS FOR MODULATION SPACES

ÁRPÁD BÉNYI, LOUKAS GRAFAKOS\*, KARLHEINZ GRÖCHENIG, AND KASSO OKOUDJOU

ABSTRACT. We prove the boundedness of a general class of Fourier multipliers, in particular of the Hilbert transform, on modulation spaces. In general, however, the Fourier multipliers in this class fail to be bounded on  $L^p$  spaces. The main tools are Gabor frames and methods from time-frequency analysis.

## 1. INTRODUCTION

In this note we explore the boundedness properties of certain translation invariant operators (initially defined on the class of Schwartz rapidly decreasing smooth functions). Namely, if  $b > 0$  and  $\mathbf{c} = (c_n)_{n \in \mathbb{Z}}$  is a bounded sequence of complex numbers, we are interested in the operator  $H_{b,\mathbf{c}}$  (formally) defined by

$$(1) \quad H_{b,\mathbf{c}} = \sum_{n=-\infty}^{+\infty} c_n (M_{bn} H M_{-bn} - M_{b(n+1)} H M_{-b(n+1)}).$$

Here,  $M_b$  denotes the modulation by  $b$  and  $H$  is the Hilbert transform, that is,

$$M_b f(t) = e^{2\pi i t \cdot b} f(t)$$

and

$$H f(x) = \frac{1}{\pi} \lim_{\epsilon \rightarrow 0} \int_{|t| > \epsilon} \frac{f(x-t)}{t} dt.$$

The operators  $H_{b,\mathbf{c}}$  are better understood if viewed on the Fourier side as multiplier operators

$$(2) \quad \widehat{H_{b,\mathbf{c}} f} = m_{b,\mathbf{c}} \widehat{f},$$

with Fourier multipliers

$$(3) \quad m_{b,\mathbf{c}} = -2i \sum_{n=-\infty}^{+\infty} c_n \chi_{(bn, b(n+1))};$$

$\chi_{(a,b)}$  denotes the characteristic function of the real interval  $(a, b)$ . It is easy to see that the Hilbert transform is a particular case of operator  $\frac{1}{2} H_{b,\mathbf{c}}$ . Indeed, if we recall

---

*Date:* January 11, 2005.

*2000 Mathematics Subject Classification.* Primary 42A45; Secondary 42B35, 42C99, 47G30.

*Key words and phrases.* Fourier multipliers, Hilbert transform, Gabor frames, modulation spaces, short-time Fourier transform.

\*Research partially supported by the NSF.

that  $m(\xi) = -isgn \xi$  is the multiplier of  $H$ , then  $H = \frac{1}{2}H_{b,\mathbf{c}}$  for any  $b > 0$  and  $\mathbf{c} = (c_n)_{n \in \mathbb{Z}}$ , with  $c_n = 1$  for  $n \geq 0$  and  $c_n = -1$  for  $n < 0$ .

There are many proofs for the boundedness of  $H$  on  $L^p$  spaces,  $1 < p < \infty$ . For example, in the context of Calderón-Zygmund theory, the boundedness of  $H$  follows from its boundedness on  $L^2$  (a consequence of Plancherel's theorem), a weak type  $(1,1)$  estimate, and interpolation. A natural question then is whether the more general operators  $H_{b,\mathbf{c}}$  are also bounded on the Lebesgue spaces  $L^p$ ,  $1 < p < \infty$ . We note right away that the boundedness of the sequence  $\mathbf{c}$  and Plancherel's theorem guarantee that  $H_{b,\mathbf{c}}$  maps  $L^2$  into  $L^2$ . But unlike the kernel of the Hilbert transform, the convolution kernels of the operators  $H_{b,\mathbf{c}}$  do not possess the required amount of smoothness to fall under the scope of Calderón-Zygmund theory. Moreover, the multipliers  $m_{b,\mathbf{c}}$  are not of bounded variation in general, nor do they satisfy the conditions of some Fourier multiplier theorem. In fact, the operators  $H_{b,\mathbf{c}}$  fail to be bounded on  $L^p$ ,  $p \neq 2$ .

Consider for example the operators  $H_{1,\mathbf{c}}$ ,  $\mathbf{c} = (c_n)_{n \in \mathbb{Z}} \in \ell^\infty$ , and assume that an estimate of the form

$$(4) \quad \|H_{1,\mathbf{c}}f\|_p \leq C\|\mathbf{c}\|_{\ell^\infty} \|f\|_p$$

were true for some  $1 < p < 2$  and a constant  $C$  independent of  $\mathbf{c}$  and  $f$ . This inequality remains true, if we replace  $c_n$  by  $\pm c_n$  for any choice of sign; in particular we may replace  $c_n$  with  $c_n r_n(t)$ , where  $r_n(t)$  is the  $n$ th Rademacher function (see [2, p. 177] for the definition of these functions.) Define the operator  $S_n$  by  $\widehat{S_n f} = c_n \chi_{[n,n+1]} \hat{f}$ . Raising (4) to the power  $p$ , integrating over  $t \in [0, 1]$ , and using a key inequality for the Rademacher functions ([9, Appendix C.2]), we deduce that for every  $f \in L^p$

$$(5) \quad \left\| \left( \sum_{n \in \mathbb{Z}} |S_n f|^2 \right)^{1/2} \right\|_{L^p} \leq C \|\mathbf{c}\|_{\ell^\infty} \|f\|_{L^p}.$$

For the choice  $c_n = 1$  this averaging procedure over the plus and minus signs yields a square function for which estimate (5) is known to hold true only when  $p \geq 2$ ; see e.g. [9, Sec. 10.2]. This contradicts our initial assumption  $1 < p < 2$ . By duality we can also exclude the case  $p > 2$ , because the space of Fourier multipliers on  $L^p$  coincides with those on  $L^{p'}$ , where  $p' = p/(p-1)$  is the dual exponent of  $p$ . We conclude that an estimate of the form (4) holds true if and only if  $p = 2$ .

Nevertheless, we will prove below that the operators  $H_{b,\mathbf{c}}$  are bounded on a different class of function spaces, the so-called modulation spaces  $\mathcal{M}^{p,q}$ ,  $1 < p < \infty$ ,  $1 \leq q \leq \infty$ . These spaces include  $L^2 = \mathcal{M}^{2,2}$  and are defined by their phase-space distribution (instead of their Littlewood-Paley decomposition). The modulation spaces occur naturally in time-frequency analysis (or phase-space analysis) and have found numerous applications to pseudodifferential operators, signal analysis, non-linear approximation, and the formulation of uncertainty principles; see, e.g., [8], [11], [12], [14], [15], [17], [18], [19], [23].

We will prove the boundedness of the Fourier multipliers by studying their matrix with respect to a so-called Gabor frame (often called Weyl-Heisenberg frame) and by using the Gabor expansion of functions. In this manner we can convert the question of boundedness of Fourier multipliers on  $\mathcal{M}^{p,q}$  into a problem about the boundedness of

an infinite matrix acting on certain sequence spaces. It is perhaps not too surprising that this discretization leads to the discrete Hilbert transform that was first studied by Hilbert in  $\ell^2$  setting and by M. Riesz [21] and Titchmarsh [22] on general  $\ell^p$  spaces.

In particular, we obtain the boundedness of the Hilbert transform on modulation spaces. This fact was first observed by Okoudjou in [20]. In this paper we treat a natural extension and also close a gap in the original proof. We also point out that a similar class of (pointwise) multipliers was considered on the so-called amalgam spaces in [24]. Since the modulation spaces are the Fourier transforms of certain amalgam spaces, some of the results in [24, Theorem 3.6] overlap with our main result. However, our techniques and the ones used in [24] are completely different. In addition, the use of time-frequency techniques in “hard analysis” seems of independent interest. As for the endpoint modulation spaces, the situation is similar to the case of Lebesgue spaces. For example, the Hilbert transform fails to be bounded on the Feichtinger algebra  $\mathcal{M}^1$ .

The remainder of the paper is organized as follows. The next section is devoted to some basic facts about modulation spaces and Gabor frames. The main result is stated and proved in Section 3, as well as some counter-example.

## 2. PRELIMINARIES

**2.1. General notation.** We will be working on the real line  $\mathbb{R}$ . The operators of translation and modulation of a function  $f$  with domain  $\mathbb{R}$  are defined by

$$T_x f(t) = f(t - x) \quad \text{and} \quad M_y f(t) = e^{2\pi i y t} f(t).$$

The Fourier transform of  $f \in L^1(\mathbb{R})$  is  $\hat{f}(\omega) = \int_{\mathbb{R}} f(t) e^{-2\pi i t \omega} dt$ ,  $\omega \in \mathbb{R}$ . The Fourier transform is an isomorphism of the Schwartz space  $\mathcal{S} = \mathcal{S}(\mathbb{R})$  onto itself, and extends to the space  $\mathcal{S}' = \mathcal{S}'(\mathbb{R})$  of tempered distributions by duality.

The inner product of two functions  $f, g \in L^2$  is  $\langle f, g \rangle = \int_{\mathbb{R}} f(t) \overline{g(t)} dt$ , and its extension to  $\mathcal{S}' \times \mathcal{S}$  will be also denoted by  $\langle \cdot, \cdot \rangle$ .

The Short-Time Fourier Transform (STFT) of a function  $f$  with respect to a window  $g$  is

$$V_g f(x, y) = \langle f, M_y T_x g \rangle = \int_{\mathbb{R}} e^{-2\pi i y t} \overline{g(t - x)} f(t) dt,$$

whenever the integral makes sense. This definition can be extended to  $f \in \mathcal{S}'(\mathbb{R})$  and  $g \in \mathcal{S}(\mathbb{R})$  and yields a continuous function  $V_g f$  of polynomial growth [10]. In a less obvious way, the STFT makes sense even in the space of tempered distributions  $\mathcal{S}'(\mathbb{R}^2)$  when both  $f \in \mathcal{S}'(\mathbb{R})$  and  $g \in \mathcal{S}'(\mathbb{R})$ , see, e.g., Folland’s book [7, Prop. 1.42].

We let  $L^{p,q} = L^{p,q}(\mathbb{R} \times \mathbb{R})$  be the spaces of measurable functions  $f(x, y)$  for which the mixed norm

$$\|f\|_{L^{p,q}} = \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |f(x, y)|^p dx \right)^{q/p} dy \right)^{1/q}$$

is finite. If  $p = q$ , we have  $L^{p,p} = L^p$ , the usual Lebesgue spaces. By  $\ell^{p,q}(\mathbb{Z} \times \mathbb{Z})$  we denote the spaces of sequences  $\mathbf{a} = (a_{k,l})_{k,l \in \mathbb{Z}}$  for which the mixed norm

$$\|\mathbf{a}\|_{\ell^{p,q}} = \left( \sum_{l \in \mathbb{Z}} \left( \sum_{k \in \mathbb{Z}} |a_{k,l}|^p \right)^{q/p} \right)^{1/q}$$

is finite. If  $p = q$ , we recover the sequence spaces  $\ell^p(\mathbb{Z} \times \mathbb{Z})$ .

## 2.2. Modulation spaces.

**Definition 1.** Given  $1 \leq p, q \leq \infty$ , and given a window function  $g \in \mathcal{S}$ , the modulation space  $\mathcal{M}^{p,q} = \mathcal{M}^{p,q}(\mathbb{R})$  is the space of all distributions  $f \in \mathcal{S}'$  for which the following norm is finite:

$$(6) \quad \|f\|_{\mathcal{M}^{p,q}} = \left( \int_{\mathbb{R}} \left( \int_{\mathbb{R}} |V_g f(x, y)|^p dx \right)^{q/p} dy \right)^{1/q} = \|V_g f\|_{L^{p,q}},$$

with the usual modifications if  $p$  and/or  $q$  are infinite. When  $p = q$ , we will write  $\mathcal{M}^p$  for the modulation space  $\mathcal{M}^{p,p}$ .

*Remark 1.* The definition is independent of the choice of the window  $g$  in the sense of equivalent norms. If  $1 \leq p, q < \infty$ , then  $\mathcal{M}^1$  is densely embedded into  $\mathcal{M}^{p,q}$ . In fact, the Schwartz class  $\mathcal{S}$  is dense in  $\mathcal{M}^{p,q}$  for  $1 \leq p, q < \infty$ . One can also show that the dual of  $\mathcal{M}^{p,q}$  is  $\mathcal{M}^{p',q'}$ , where  $1 \leq p, q < \infty$  and  $\frac{1}{p} + \frac{1}{p'} = \frac{1}{q} + \frac{1}{q'} = 1$ . We refer to [10] and the references therein for more details about modulation spaces.

*Remark 2.* The modulation space  $\mathcal{M}^1$ , also called the Feichtinger algebra, is a Banach algebra under both pointwise multiplication and convolution and is invariant under Fourier transform. It plays also an important role in the theory of Gabor frames where it serves as a convenient class of windows that generate Gabor frames for the whole class of modulation spaces.

## 2.3. Gabor Frames.

**Definition 2.** Given a window function  $\phi \in L^2(\mathbb{R})$  and constants  $\alpha, \beta > 0$ , we say that  $\{M_{\beta n} T_{\alpha k} \phi\}_{k,n \in \mathbb{Z}}$  is a Gabor frame for  $L^2(\mathbb{R})$  if there exist constants  $A, B > 0$  (called frame bounds) such that

$$A \|f\|_{L^2(\mathbb{R})}^2 \leq \sum_{k,n \in \mathbb{Z}} |\langle f, M_{\beta n} T_{\alpha k} \phi \rangle|^2 \leq B \|f\|_{L^2(\mathbb{R})}^2, \quad \forall f \in L^2(\mathbb{R}).$$

We refer to the works by Daubechies [1], Gröchenig [10], and Heil and Walnut [16] for extensive treatments of frames and Gabor frames.

The theory of Gabor frames can be generalized from the pure  $L^2$ -theory to the whole class of modulation spaces. The next theorem taken from [5], [6], and [10] provides a characterization of modulation spaces by means of Gabor frames and will be used heavily in the sequel.

**Theorem A.** *Let  $\phi \in \mathcal{M}^1$  be such that  $\{M_{\beta n} T_{\alpha k} \phi\}_{k,n \in \mathbb{Z}}$  is a Gabor frame for  $L^2$ , and let  $1 \leq p, q \leq \infty$ . Then there exists a (canonical) dual  $\gamma \in \mathcal{M}^1$  such that every*

tempered distribution in  $\mathcal{M}^{p,q}$  has a Gabor expansion that converges unconditionally (or weak\* unconditionally if  $p = \infty$  or  $q = \infty$ ), namely

$$f = \sum_{k,n \in \mathbb{Z}} \langle f, M_{\beta n} T_{\alpha k} \gamma \rangle M_{\beta n} T_{\alpha k} \phi, \quad \forall f \in \mathcal{M}^{p,q}(\mathbb{R});$$

moreover, we have the following norm equivalences

$$\|f\|_{\mathcal{M}^{p,q}} \asymp \|\langle f, M_{\beta n} T_{\alpha k} \phi \rangle\|_{\ell^{p,q}} \asymp \|\langle f, M_{\beta n} T_{\alpha k} \gamma \rangle\|_{\ell^{p,q}}.$$

To summarize, a tempered distribution  $f$  belongs to the modulation space  $\mathcal{M}^{p,q}(\mathbb{R})$  if and only if the sequence of its Gabor coefficients defined as

$$C_\phi f = (\langle f, M_{\beta n} T_{\alpha k} \phi \rangle)_{k,n \in \mathbb{Z}}$$

belongs to the sequence space  $\ell^{p,q}(\mathbb{Z} \times \mathbb{Z})$ . Moreover, the norm of  $f$  is equivalent to the norm of its Gabor coefficients.

### 3. BOUNDEDNESS OF $H_{b,\mathbf{c}}$ ON MODULATION SPACES

Our main result can be stated as follows:

**Theorem 1.** *For any  $b > 0$  and  $\mathbf{c} \in \ell^\infty$ , the operators  $H_{b,\mathbf{c}}$  are bounded from  $\mathcal{M}^{p,q}$  into  $\mathcal{M}^{p,q}$  for  $1 < p < \infty$ ,  $1 \leq q \leq \infty$  with a norm estimate*

$$\|H_{b,\mathbf{c}} f\|_{\mathcal{M}^{p,q}} \leq C \|\mathbf{c}\|_\infty \|f\|_{\mathcal{M}^{p,q}}$$

for some constant depending only on  $b, p$ , and  $q$ . In particular, the Hilbert transform  $H$  is bounded on  $\mathcal{M}^{p,q}$  for  $1 < p < \infty$  and  $1 \leq q \leq \infty$

*Proof.* Since the modulation spaces are invariant under dilations, we may assume without loss of generality that  $b = 1$  by conjugating  $H_{b,\mathbf{c}}$  with a suitable dilation. Thus from now on we will only consider the multiplier (1) with  $b = 1$ .

Next we choose a Gabor frame that is tailored to the analysis of our particular class of Fourier multipliers. Let

$$\phi(x) = \left(\frac{\sin \pi x}{\pi x}\right)^2, \text{ or equivalently, } \hat{\phi}(\omega) = \chi_{[-1/2, 1/2]} * \chi_{[-1/2, 1/2]}(\omega) = \max(0, 1 - |\omega|).$$

It follows that  $\phi \in \mathcal{M}^1$  [3] or [10, Prop. 12.1.6], and that  $\{M_n T_{\frac{k}{2}} \phi\}_{k,n \in \mathbb{Z}}$  is a Gabor frame for  $L^2(\mathbb{R})$  [10, Thm. 6.4.1].

We expand  $f$  with respect to a Gabor frame, i.e.,  $f = \sum_{k,n \in \mathbb{Z}} \langle f, M_{\beta n} T_{\alpha k} \gamma \rangle M_{\beta n} T_{\alpha k} \phi$ , and then take the coefficients of  $H_{1,\mathbf{c}}$ . We find that

$$\begin{aligned} (7) \quad (C_\phi H_{1,\mathbf{c}} f)(k, n) &= \langle H_{1,\mathbf{c}} f, M_n T_{\frac{k}{2}} \phi \rangle \\ &= \sum_{k', n' \in \mathbb{Z}} \langle f, M_{n'} T_{\frac{k'}{2}} \gamma \rangle \langle H_{1,\mathbf{c}} M_{n'} T_{\frac{k'}{2}} \phi, M_n T_{\frac{k}{2}} \phi \rangle. \end{aligned}$$

Recall that by Theorem A,  $f \in \mathcal{M}^{p,q}$  if and only if  $C_\gamma f = (\langle f, M_n T_{\frac{k}{2}} \gamma \rangle)_{k,n \in \mathbb{Z}} \in \ell^{p,q}(\mathbb{Z}^2)$  and  $H_{1,\mathbf{c}} f \in \mathcal{M}^{p,q}$  if and only if  $C_\phi H_{1,\mathbf{c}} f = (\langle H_{1,\mathbf{c}} f, M_n T_{\frac{k}{2}} \phi \rangle)_{k,n \in \mathbb{Z}} \in \ell^{p,q}(\mathbb{Z}^2)$ .

Consequently to verify the boundedness of  $H_{1,\mathbf{c}}$ , it suffices to show that the matrix  $\Upsilon$  whose entries are defined by

$$\Upsilon_{(k,n),(k',n')} = \langle H_{1,\mathbf{c}} M_{n'} T_{\frac{k'}{2}} \phi, M_n T_{\frac{k}{2}} \phi \rangle$$

is bounded on  $\ell^{p,q}(\mathbb{Z}^2)$ . The commutative diagram (8) illustrates the situation.

$$(8) \quad \begin{array}{ccc} \mathcal{M}^{p,q} & \xrightarrow{H_{1,\mathbf{c}}} & \mathcal{M}^{p,q} \\ \downarrow C_\gamma & & \downarrow C_\phi \\ \ell^{p,q} & \xrightarrow{\Upsilon} & \ell^{p,q} \end{array}$$

We now compute the entries of the matrix  $\Upsilon$ . Here we exploit the special properties of the basic function  $\phi$  and the algebra of time-frequency shifts. We have:

$$(9) \quad \begin{aligned} \Upsilon_{(k,n),(k',n')} &= \langle H_{1,\mathbf{c}} M_{n'} T_{\frac{k'}{2}} \phi, M_n T_{\frac{k}{2}} \phi \rangle \\ &= \langle (H_{1,\mathbf{c}} M_{n'} T_{\frac{k'}{2}} \phi)^\wedge, (M_n T_{\frac{k}{2}} \phi)^\wedge \rangle \\ &= \left\langle \sum_{m \in \mathbb{Z}} c_m \chi_{[m,m+1]} \cdot T_{n'} M_{-\frac{k'}{2}} \hat{\phi}, T_n M_{-\frac{k}{2}} \hat{\phi} \right\rangle \\ &= \sum_{m \in \mathbb{Z}} c_m e^{\pi i k' (n' - n)} \langle (T_{-n} \chi_{[m,m+1]}) M_{\frac{k-k'}{2}} T_{n'-n} \hat{\phi}, \hat{\phi} \rangle \\ &= \sum_{m \in \mathbb{Z}} c_m (-1)^{k'(n'-n)} \langle M_{\frac{k-k'}{2}} \chi_{[m-n, m-n+1]} T_{n'-n} \hat{\phi}, \hat{\phi} \rangle. \end{aligned}$$

Because the function  $\hat{\phi}$  is compactly supported on  $[-1, 1]$ , the sum in (9) is finite, and only the terms  $m = n - 1$  and  $m = n$  occur:

$$(10) \quad \begin{aligned} \Upsilon_{(k,n),(k',n')} &= c_{n-1} (-1)^{k'(n'-n)} \langle M_{\frac{k-k'}{2}} \chi_{[-1,0]} T_{n'-n} \hat{\phi}, \hat{\phi} \rangle \\ &\quad + c_n (-1)^{k'(n'-n)} \langle M_{\frac{k-k'}{2}} \chi_{[0,1]} T_{n'-n} \hat{\phi}, \hat{\phi} \rangle. \end{aligned}$$

Furthermore, the support condition on  $\hat{\phi}$  implies that the indices  $n, n'$  are related by  $n - 1 \leq n' \leq n + 1$  and we only need to calculate the following integrals explicitly:

$$\begin{aligned} \langle M_{\frac{l}{2}} \chi_{[0,1]} \hat{\phi}, \hat{\phi} \rangle &= \langle M_{-\frac{l}{2}} \chi_{[-1,0]} \hat{\phi}, \hat{\phi} \rangle = \int_0^1 (1 - \omega)^2 e^{\pi i l \omega} d\omega, \\ \langle M_{\frac{l}{2}} \chi_{[0,1]} T_1 \hat{\phi}, \hat{\phi} \rangle &= \langle M_{-\frac{l}{2}} \chi_{[-1,0]} T_{-1} \hat{\phi}, \hat{\phi} \rangle = \int_0^1 \omega (1 - \omega) e^{\pi i l \omega} d\omega, \end{aligned}$$

whereas

$$\langle M_{\frac{l}{2}} \chi_{[0,1]} T_{-1} \hat{\phi}, \hat{\phi} \rangle = \langle M_{-\frac{l}{2}} \chi_{[-1,0]} T_1 \hat{\phi}, \hat{\phi} \rangle = 0.$$

The evaluation of these elementary integrals (with integration by parts) yields  $\int_0^1 (1 - \omega)^2 e^{\pi i l \omega} d\omega = \rho_l + \delta_l/3$  and  $\int_0^1 \omega (1 - \omega) e^{\pi i l \omega} d\omega = \epsilon_l + \delta_l/6$ , where

$$\delta_k = \begin{cases} 0 & \text{if } k \neq 0 \\ 1 & \text{if } k = 0, \end{cases}$$

$$\rho_k = \begin{cases} -\frac{1}{\pi i k} + \frac{2}{\pi^2 k^2} + \frac{2((-1)^k - 1)}{(i\pi k)^3} & \text{if } k \neq 0 \\ 0 & \text{if } k = 0 \end{cases}$$

and

$$\epsilon_k = \begin{cases} -\frac{(-1)^{k+1}}{\pi^2 k^2} + \frac{2((-1)^k - 1)}{(i\pi k)^3} & \text{if } k \neq 0 \\ 0 & \text{if } k = 0. \end{cases}$$

Using these sequences we can compute the entries of  $\Lambda$  as follows:

- Case 1: If  $n' = n$ , then

$$\Upsilon_{(k,n),(k',n)} = c_{n-1} (\rho_{k'-k} + \frac{1}{3}\delta_{k-k'}) + c_n (\rho_{k-k'} + \frac{1}{3}\delta_{k'-k}).$$

- Case 2: If  $n' = n + 1$ , then

$$\Upsilon_{(k,n),(k',n')} = c_n (-1)^{k'} (\epsilon_{k-k'} + \frac{1}{6}\delta_{k-k'}).$$

- Case 3: If  $n' = n - 1$ , then

$$\Upsilon_{(k,n),(k',n')} = c_{n-1} (-1)^{k'} (\epsilon_{k'-k} + \frac{1}{6}\delta_{k'-k}).$$

Consequently, letting  $a_{k',n'} = \langle f, M_{n'} T_{\frac{k'}{2}} \gamma \rangle$ , we can rewrite (7) as follows:

$$\begin{aligned} (C_\phi H_{1,c}(f))(k, n) &= \sum_{k', n' \in \mathbb{Z}} \Upsilon_{(k,n),(k',n')} a_{k',n'} \\ &= \sum_{k' \in \mathbb{Z}} a_{k',n} c_n (\rho_{k-k'} + \frac{1}{3}\delta_{k'-k}) + \sum_{k' \in \mathbb{Z}} a_{k',n} c_{n-1} (\rho_{k'-k} + \frac{1}{3}\delta_{k-k'}) \\ &\quad + \sum_{k' \in \mathbb{Z}} a_{k',n+1} c_n (-1)^{k'} (\epsilon_{k-k'} + \frac{1}{6}\delta_{k-k'}) \\ (11) \quad &\quad + \sum_{k' \in \mathbb{Z}} a_{k',n-1} c_{n-1} (-1)^{k'} (\epsilon_{k'-k} + \frac{1}{6}\delta_{k'-k}). \end{aligned}$$

The action of the matrix  $\Upsilon$  can now be expressed in terms of convolutions with the sequences  $\rho$ ,  $\epsilon$  and  $\delta$  defined above. More precisely,

$$\begin{aligned} (C_\phi H_{1,c}(f))(k, n) &= c_n (a_{\cdot,n} * \rho(k) + \frac{1}{3}a_{k,n}) + c_{n-1} (a_{\cdot,n} * \tilde{\rho}(k) + \frac{1}{3}a_{k,n}) \\ &\quad + c_n ((-1) \cdot a_{\cdot,n+1} * \epsilon(k) + \frac{1}{6}a_{k,n+1}) \\ (12) \quad &\quad + c_{n-1} ((-1) \cdot a_{\cdot,n-1} * \tilde{\epsilon}(k) + \frac{1}{6}a_{k,n-1}), \end{aligned}$$

where  $\tilde{x}(k) = x(-k)$  and  $(-1) \cdot a_{\cdot,n} * \epsilon$  denotes the sequence  $\left( \sum_{k' \in \mathbb{Z}} (-1)^{k'} a_{k',n} \epsilon_{k-k'} \right)_{k \in \mathbb{Z}}$

for any fixed  $n$ .

The sequence  $\epsilon$  belongs to  $\ell^1(\mathbb{Z})$ , thus we can use Young's inequality to take care of the convolution terms involving  $\epsilon$ . However,  $\rho$  is the sum of three sequences, two of which are in  $\ell^1(\mathbb{Z})$ , but the third one,  $\frac{1}{\pi ik}$ , is clearly not summable. Fortunately, the convolution with  $\frac{1}{\pi ik}$  (the discrete Hilbert transform) is bounded on  $\ell^p(\mathbb{Z})$  for  $1 < p < \infty$ ; see the book by Hardy, Littlewood and Pólya [13, Section 8.12], also [21], [22]. Consequently, convolution with  $\rho$  is also bounded on  $\ell^p(\mathbb{Z})$  for  $1 < p < \infty$ , and using Young's inequality we obtain

$$\left(\sum_{k \in \mathbb{Z}} |C_\phi H_{1,c}(f)(k, n)|^p\right)^{1/p} \leq K \|\mathbf{c}\|_\infty \|a_{\cdot, n}\|_p$$

for some positive constant  $K$  and every  $n \in \mathbb{Z}$ . Hence, by taking the  $q$ -norm with respect to the variable  $n$  we obtain

$$\|C_\phi H_{1,c}(f)\|_{\ell^{p,q}(\mathbb{Z}^2)} \leq K \|\mathbf{c}\|_\infty \|a\|_{\ell^{p,q}(\mathbb{Z}^2)},$$

for all  $1 \leq q \leq \infty$ . This concludes the proof.  $\square$

*Remark 3.* In general, the operators  $H_{b,c}$  are not bounded on  $\mathcal{M}^1$ . In fact, this is already the case for the Hilbert transform  $H$ . Assume on the contrary that  $Hf \in \mathcal{M}^1$  for every  $f \in \mathcal{M}^1$ . Since  $\mathcal{M}^1$  is invariant under the Fourier transform, this would imply that  $\widehat{Hf} = -i \operatorname{sgn}(\cdot) \widehat{f} \in \mathcal{M}^1$  and  $\widehat{Hf}$  would have a discontinuity at the origin whenever  $\widehat{f}(0) \neq 0$ . But this contradicts the fact that every function in  $\mathcal{M}^1$  is uniformly continuous. By duality,  $H$  cannot be bounded from  $\mathcal{M}^\infty$  into  $\mathcal{M}^\infty$  either.

#### 4. ACKNOWLEDGMENT

The authors thank Chris Heil and Norbert Kaiblinger for very helpful and interesting discussions while completing this work.

#### REFERENCES

- [1] I. Daubechies, "Ten Lectures on Wavelets," SIAM, Philadelphia PA, 1992.
- [2] J. Duandikoetxea, "Fourier Analysis," Graduate Studies in Mathematics **29**, AMS, Providence RI, 2001.
- [3] H. G. Feichtinger, *On a new Segal algebra*, Monatsh. Math. **92** (1981), 269–289.
- [4] H. G. Feichtinger, *Modulation spaces on locally Abelian groups*, Technical Report, University of Vienna, 1983, Updated version appeared in Proceedings of "International Conference on Wavelets and Applications" 2002, pp. 99-140, Chennai, India, 2003.
- [5] H. G. Feichtinger and K. Gröchenig, *Banach spaces related to integrable group representations and their atomic decompositions I*, J. Funct. Anal. **86** (1989), 307–334.
- [6] H. G. Feichtinger and K. Gröchenig, *Banach spaces related to integrable group representations and their atomic decompositions II*, Monatsh. Math. **108** (1989), 129–148.
- [7] G. B. Folland, "Harmonic Analysis in Phase Space," Ann. of Math. Studies, Princeton University Press, Princeton NJ, 1989.
- [8] E. Galperin and K. Gröchenig, *Uncertainty principles as embeddings of modulation spaces*, J. Math. Anal. Appl. **271** (2002), 181–202.
- [9] L. Grafakos, "Classical and Modern Fourier Analysis," Prentice Hall, Upper Saddle River NJ, 2003.
- [10] K. Gröchenig, "Foundations of Time-Frequency Analysis," Birkhäuser, Boston MA, 2001.



- [11] K. Gröchenig and C. Heil, *Modulation spaces and pseudodifferential operators*, Int. Eq. Oper. Theory **34** (1999), 439–457.
- [12] K. Gröchenig and C. Heil, *Counterexamples for boundedness of pseudodifferential operators*, Osaka J. Math. **41** (2004), 1–11.
- [13] G. H. Hardy, J. E. Littlewood and G. Pólya, “Inequalities,” Cambridge University Press, Cambridge, 1952.
- [14] C. Heil: *Integral operators, pseudodifferential operators, and Gabor frames*, in: “Advances in Gabor Analysis,” H. G. Feichtinger and T. Strohmer, eds., Birkhäuser, Boston (2003), 153–169.
- [15] C. Heil, J. Ramanathan and P. Topiwala, *Singular values of compact pseudodifferential operators*, J. Funct. Anal. **150** (1997), 426–452.
- [16] C. Heil and D. Walnut, *Continuous and discrete wavelet transforms*, SIAM Review **31** (1989), 628–666.
- [17] J. A. Hogan and J. D. Lakey, *Embeddings and uncertainty principles for generalized modulation spaces*, in: “Modern Sampling Theory,” J. J. Benedetto and P. J. S. G. Ferreira, eds., Birkhäuser, Boston (2001), 73–105.
- [18] D. Labate, *Time-frequency analysis of pseudodifferential operators*, Monatsh. Math. **133** (2001), 143–156.
- [19] D. Labate, *Pseudodifferential operators on modulation spaces*, J. Math. Anal. Appl. **262** (2001), 242–255.
- [20] K. A. Okoudjou, *Characterization of function spaces and boundedness of bilinear pseudodifferential operators through Gabor frames*, Ph.D. thesis, Georgia Institute of Technology, Atlanta, 2003.
- [21] M. Riesz, *Sur les fonctions conjuguées*, Math. Zeit. **27** (1928), 218–244.
- [22] E. C. Titchmarsh, *An inequality in the theory of series*, J. London Math. Soc. **3** (1928), 81–83.
- [23] J. Toft, *Modulation spaces and pseudodifferential operators I*, J. Funct. Anal. **207** (2004), 399–429.
- [24] M. L. Torres de Squire, *Multipliers for amalgams and the algebra  $S_0(G)$* , Can. J. Math., **39**(1) (1987), 123–148.

ÁRPÁD BÉNYI, DEPARTMENT OF MATHEMATICS AND STATISTICS, LEDERLE GRT, UNIVERSITY OF MASSACHUSETTS, AMHERST, MA 01003, USA

*E-mail address:* benyi@math.umass.edu

LOUKAS GRAFAKOS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA

*E-mail address:* loukas@math.missouri.edu

KARLHEINZ GRÖCHENIG, GSF RESEARCH CENTER, DEPARTMENT OF BIOMATHEMATICS AND BIOMETRY, INGOLSTÄDTER STR. 1, D-85764 NEUHERBERG, GERMANY

*E-mail address:* karlheinz.groechenig@gsf.de

KASSO OKOUDJOU, DEPARTMENT OF MATHEMATICS, MALOTT HALL, CORNELL UNIVERSITY, ITHACA, NY 14853, USA

*E-mail address:* kasso@math.cornell.edu