

THE MARCINKIEWICZ MULTIPLIER THEOREM REVISITED

LOUKAS GRAFAKOS AND LENKA SLAVÍKOVÁ

ABSTRACT. We provide a complete proof of an optimal version of the Marcinkiewicz multiplier theorem and discuss a relevant example.

1. INTRODUCTION AND STATEMENT OF RESULTS

We revisit a product-type Sobolev space version of the Marcinkiewicz multiplier theorem. A version of this result first appeared in Carbery [2] but a stronger version of it is a consequence of the work of Carbery and Seeger [3]. In this note we provide a self-contained proof of the Marcinkiewicz multiplier theorem, we point out that the conditions on the indices are optimal, and we provide a comparison with the Hörmander multiplier theorem, which indicates that the former is stronger than the latter.

Given a bounded function σ on \mathbb{R}^n , we define a linear operator

$$T_\sigma(f)(x) = \int_{\mathbb{R}^n} \widehat{f}(\xi)\sigma(\xi)e^{2\pi i x \cdot \xi} d\xi$$

acting on Schwartz functions f on \mathbb{R}^n ; here $\widehat{f}(\xi) = \int_{\mathbb{R}^n} f(x)e^{-2\pi i x \cdot \xi} dx$ is the Fourier transform of f . An old problem in harmonic analysis is to find optimal sufficient conditions on σ so that the operator T_σ admits a bounded extension from $L^p(\mathbb{R}^n)$ to itself for a given $p \in (1, \infty)$. If this is the case for a given σ , then we say that σ is an L^p Fourier multiplier.

We recall the classical Marcinkiewicz multiplier theorem: Let $I_j = (-2^{j+1}, -2^j] \cup [2^j, 2^{j+1})$ for $j \in \mathbb{Z}$. Let σ be a bounded function on \mathbb{R}^n such that for all $\alpha = (\alpha_1, \dots, \alpha_n)$ with $|\alpha_1|, \dots, |\alpha_n| \leq 1$ the derivatives $\partial^\alpha \sigma$ are continuous up to the boundary of any rectangle $I_{j_1} \times \dots \times I_{j_n}$ on \mathbb{R}^n . Assume that there is a constant $A < \infty$ such that for all partitions $\{s_1, \dots, s_k\} \cup \{r_1, \dots, r_\ell\} = \{1, 2, \dots, n\}$ with $n = k + \ell$ and all $\xi_{j_i} \in I_{j_i}$ we

Mathematics Subject Classification: Primary 42B15. Secondary 42B25.

Keywords and phrases: Multiplier theorems, Sobolev spaces, interpolation.

The first author acknowledges the support of the Simons Foundation and of the University of Missouri Research Board.

have

$$(1.1) \quad \sup_{\xi_{r_1} \in I_{j_{r_1}}} \cdots \sup_{\xi_{r_\ell} \in I_{j_{r_\ell}}} \int_{I_{j_{s_1}}} \cdots \int_{I_{j_{s_k}}} |(\partial_{s_1} \cdots \partial_{s_k} \sigma)(\xi_1, \dots, \xi_n)| d\xi_{s_k} \cdots d\xi_{s_1} \leq A$$

for all $(j_1, \dots, j_n) \in \mathbb{Z}^n$. Then σ is an L^p Fourier multiplier on \mathbb{R}^n whenever $1 < p < \infty$, and there is a constant $C_{p,n} < \infty$ such that

$$(1.2) \quad \|T_\sigma\|_{L^p \rightarrow L^p} \leq C_{p,n} (A + \|\sigma\|_{L^\infty}).$$

To obtain a Sobolev space version of this result we define $(I - \partial_\ell^2)^{\frac{\gamma_\ell}{2}} f$ as the linear operator $((1 + 4\pi^2|\eta_\ell|^2)^{\frac{\gamma_\ell}{2}} \widehat{f}(\eta))^\vee$ associated with the multiplier $(1 + 4\pi^2|\eta_\ell|^2)^{\frac{\gamma_\ell}{2}}$. The purpose of this note is to provide a self-contained exposition of a version of the Marcinkiewicz multiplier theorem which requires only $1/r + \varepsilon$ derivatives per variable in L^r instead of a full derivative in L^1 as in (1.1).

Theorem 1.1. *Let $n \in \mathbb{N}$, $n \geq 2$. Suppose that $1 \leq r < \infty$ and ψ is a Schwartz function on the line whose Fourier transform is supported in $[-2, -1/2] \cup [1/2, 2]$ and which satisfies $\sum_{j \in \mathbb{Z}} \widehat{\psi}(2^{-j}\xi) = 1$ for all $\xi \neq 0$. Let $\gamma_\ell > 1/r$, $\ell = 1, \dots, n$. If a function σ on \mathbb{R}^n satisfies*

$$(1.3) \quad \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} (\widehat{\psi}(\xi_1) \cdots \widehat{\psi}(\xi_n) \sigma(2^{j_1}\xi_1, \dots, 2^{j_n}\xi_n)) \right\|_{L^r} < \infty,$$

then T_σ admits a bounded extension from $L^p(\mathbb{R}^n)$ to itself for all $1 < p < \infty$ with

$$(1.4) \quad \left| \frac{1}{p} - \frac{1}{2} \right| < \min(\gamma_1, \dots, \gamma_n).$$

Moreover, (1.4) is optimal in the sense that if T_σ is L^p -bounded for every σ satisfying (1.3), then the strict inequality in (1.4) must necessarily hold.

Carbery [2] first formulated a version of Theorem 1.1 in which the multiplier lies in a product-type L^2 -based Sobolev space. Carbery and Seeger [3, Remark after Prop. 6.1] obtained Theorem 1.1 in the case when $\gamma_1 = \cdots = \gamma_n > \left| \frac{1}{p} - \frac{1}{2} \right| = \frac{1}{r}$. The positive direction of their result also appeared in [4, Condition (1.4)] but this time the range of p is $\left| \frac{1}{p} - \frac{1}{2} \right| < \frac{1}{r}$ and is expressed in terms of the integrability of the multiplier and not in terms of its smoothness. In our opinion it is more natural, nonetheless, to state condition (1.4) in terms of the smoothness of the multiplier, as in the case of the sharp version of the Hörmander multiplier theorem, see [11].

The class of multipliers which satisfies the assumptions of Theorem 1.1 is strictly larger than the set of multipliers treated by the version of the Hörmander multiplier theorem due to Calderón and Torchinsky [1, Theorem 4.6]; see also [8]. Since Theorem 1.1

assumes the multiplier σ to have $1/r+\varepsilon$ derivatives in each variable, while the Hörmander multiplier theorem requires more than n/r derivatives in all variables, it is apparent that there are multipliers which can be treated by Theorem 1.1, but not by [1, Theorem 4.6]. On the other hand, it is an easy consequence of the following theorem that every multiplier satisfying the assumptions of the Hörmander multiplier theorem falls under the scope of Theorem 1.1 as well.

Theorem 1.2. *Let ψ be a Schwartz function on the line whose Fourier transform is supported in the set $\{\xi : \frac{1}{2} \leq |\xi| \leq 2\}$ and which satisfies $\sum_{k \in \mathbb{Z}} \widehat{\psi}(2^k \xi) = 1$ for every $\xi \neq 0$. Also, let Φ be a Schwartz function on \mathbb{R}^n having analogous properties. If $1 < r < \infty$ and $\gamma_1, \dots, \gamma_n$ are real numbers larger than $\frac{1}{r}$, then*

$$(1.5) \quad \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^r} \\ \leq C \sup_{j \in \mathbb{Z}} \left\| (I - \Delta)^{\frac{\gamma_1 + \dots + \gamma_n}{2}} \left[\sigma(2^j \xi) \widehat{\Phi}(\xi) \right] \right\|_{L^r}.$$

2. THE PROOF OF THEOREM 1.1; THE MAIN ESTIMATE

We start this section by introducing some notation that will be used throughout the paper. We will use ψ to denote the bump from Theorem 1.1; further, θ will stand for the function on the line satisfying

$$\widehat{\theta}(\eta) = \widehat{\psi}(\eta/2) + \widehat{\psi}(\eta) + \widehat{\psi}(2\eta).$$

One can observe that $\widehat{\theta}$ is supported in $\{\frac{1}{4} \leq |\xi| \leq 4\}$ and $\widehat{\theta} = 1$ on the support of $\widehat{\psi}$.

To simplify the notation, if $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$ and $J = (j_1, \dots, j_n) \in \mathbb{Z}^n$, we shall write

$$2^J \xi = (2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n)$$

and

$$\widehat{\psi}(\xi) = \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell), \quad \widehat{\theta}(\xi) = \prod_{\ell=1}^n \widehat{\theta}(\xi_\ell).$$

Let $k \in 1, \dots, n$. For $j \in \mathbb{Z}$ we define the Littlewood-Paley operators associated to the bumps ψ and θ by

$$\Delta_j^k f = \Delta_j^{\psi, k} f = \int_{\mathbb{R}} f(x_1, \dots, x_{k-1}, x_k - y, x_{k+1}, \dots, x_n) 2^j \psi(2^j y) dy$$

and

$$\Delta_j^{\theta, k} f = \int_{\mathbb{R}} f(x_1, \dots, x_{k-1}, x_k - y, x_{k+1}, \dots, x_n) 2^j \theta(2^j y) dy.$$

We begin with the following lemma:

Lemma 2.1. *Let $1 \leq r < \infty$, let $1 \leq \rho < 2$ satisfy $1 \leq \rho \leq r$ and let $\gamma_1, \dots, \gamma_n$ be real numbers such that $\gamma_\ell \rho > 1$, $\ell = 1, \dots, n$. Then, for any function f on \mathbb{R}^n and for all integers j_1, \dots, j_n , we have*

$$(2.6) \quad |\Delta_{j_1}^1 \cdots \Delta_{j_n}^n T_\sigma(f)| \leq C K \left[M^{(1)} \cdots M^{(n)} (|\Delta_{j_1}^{\theta,1} \cdots \Delta_{j_n}^{\theta,n} f|^\rho) \right]^{\frac{1}{\rho}},$$

where $M^{(\ell)}$ denotes the one-dimensional Hardy-Littlewood maximal operator in the ℓ -th coordinate and

$$K = \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n) \widehat{\psi}(\xi_1) \cdots \widehat{\psi}(\xi_n) \right] \right\|_{L^r}.$$

Proof. Throughout the proof we shall use the notation introduced at the beginning of this section and, whenever $J = (j_1, \dots, j_n)$, we shall write

$$\Delta_J f = \Delta_{j_1}^1 \cdots \Delta_{j_n}^n f, \quad \Delta_J^\theta f = \Delta_{j_1}^{\theta,1} \cdots \Delta_{j_n}^{\theta,n} f.$$

Since $\widehat{\theta}$ is equal to 1 on the support of $\widehat{\psi}$, we have

$$\begin{aligned} \Delta_J T_\sigma(f)(x_1, \dots, x_n) &= \int_{\mathbb{R}^n} \widehat{f}(\xi) \widehat{\psi}(2^{-J} \xi) \sigma(\xi) e^{2\pi i x \cdot \xi} d\xi \\ &= \int_{\mathbb{R}^n} \widehat{f}(\xi) \widehat{\theta}(2^{-J} \xi) \widehat{\psi}(2^{-J} \xi) \sigma(\xi) e^{2\pi i x \cdot \xi} d\xi = \int_{\mathbb{R}^n} (\Delta_J^\theta f)^\wedge(\xi) \widehat{\psi}(2^{-J} \xi) \sigma(\xi) e^{2\pi i x \cdot \xi} d\xi \\ &= \int_{\mathbb{R}^n} 2^{j_1 + \dots + j_n} (\Delta_J^\theta f)^\wedge(2^J \xi') \widehat{\psi}(\xi') \sigma(2^J \xi') e^{2\pi i (2^J x \cdot \xi')} d\xi' \\ &= 2^{j_1 + \dots + j_n} \int_{\mathbb{R}^n} (\Delta_J^\theta f)(y) \left[\widehat{\psi}(\xi') \sigma(2^J \xi') \right]^\wedge(2^J(x - y)) dy \\ &= \int_{\mathbb{R}^n} \frac{2^{j_1 + \dots + j_n} (\Delta_J^\theta f)(y)}{\prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell}} \cdot \prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell} \left[\widehat{\psi}(\xi) \sigma(2^J \xi) \right]^\wedge(2^J(x - y)) dy. \end{aligned}$$

Hölder's inequality thus yields that $|\Delta_J T_\sigma(f)(x)|$ is bounded by

$$\begin{aligned} &\left(\int_{\mathbb{R}^n} 2^{j_1 + \dots + j_n} \frac{|(\Delta_J^\theta f)(y)|^\rho}{\prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell \rho}} dy \right)^{\frac{1}{\rho}} \\ &\cdot \left(\int_{\mathbb{R}^n} 2^{j_1 + \dots + j_n} \left| \prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell} \cdot \left[\widehat{\psi}(\xi) \sigma(2^J \xi) \right]^\wedge(2^J(x - y)) \right|^{\rho'} dy \right)^{\frac{1}{\rho'}}, \end{aligned}$$

where, when $\rho = 1$, the second term in the product is to be interpreted as

$$\left\| \prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell} \cdot \left[\widehat{\psi}(\xi) \sigma(2^J \xi) \right]^\wedge(2^J(x - y)) \right\|_{L^\infty}.$$

Since $\gamma_\ell \rho > 1$ for all $\ell = 1, \dots, n$, n consecutive applications of [7, Theorem 2.1.10] yield the estimate

$$\left(\int_{\mathbb{R}^n} 2^{j_1 + \dots + j_n} \frac{|\Delta_J^\theta f(y)|^\rho}{\prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell \rho}} dy \right)^{\frac{1}{\rho}} \leq C \left[M^{(1)} \dots M^{(n)} (|\Delta_J^\theta f|^\rho)(x) \right]^{\frac{1}{\rho}}.$$

We now write

$$\begin{aligned} & \left(\int_{\mathbb{R}^n} 2^{j_1 + \dots + j_n} \left| \prod_{\ell=1}^n (1 + 2^{j_\ell} |x_\ell - y_\ell|)^{\gamma_\ell} \left[\widehat{\psi}(\xi) \sigma(2^J \xi) \right]^\wedge (2^J(x - y)) \right|^{\rho'} dy \right)^{\frac{1}{\rho'}} \\ & \leq \left(\int_{\mathbb{R}^n} \left| \prod_{\ell=1}^n (1 + |y_\ell|^2)^{\frac{\gamma_\ell}{2}} \left[\widehat{\psi}(\xi) \sigma(2^J \xi) \right]^\wedge (y) \right|^{\rho'} dy \right)^{\frac{1}{\rho'}} \\ & \leq \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \dots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^J \xi) \widehat{\psi}(\xi) \right] \right\|_{L^\rho} \\ (2.7) \quad & \leq C \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \dots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^J \xi) \widehat{\psi}(\xi) \right] \right\|_{L^r} \\ & \leq CK. \end{aligned}$$

Notice that the second inequality is the Hausdorff-Young inequality, while (2.7) is a consequence of the Kato-Ponce inequality [10] (if $\rho < r$). A combination of the preceding estimates yields (2.6). \square

Proposition 2.2. *Let $1 \leq r < \infty$ and let $\gamma_\ell > \max\{1/2, 1/r\}$, $\ell = 1, \dots, n$. If a function σ on \mathbb{R}^n satisfies (1.3), then T_σ admits a bounded extension from $L^p(\mathbb{R}^n)$ to itself for all $1 < p < \infty$.*

Proof. Suppose first that $p > 2$. Since $\gamma_\ell > \max\{1/2, 1/r\}$, $\ell = 1, \dots, n$, we can find $\rho \in [1, 2)$ such that $\rho \leq r$ and $\rho \gamma_\ell > 1$, $\ell = 1, \dots, n$. Then

$$\begin{aligned} \|T_\sigma(f)\|_{L^p(\mathbb{R}^n)} & \leq C_p(n) \left\| \left(\sum_{j_1, \dots, j_n \in \mathbb{Z}} |\Delta_{j_1}^1 \dots \Delta_{j_n}^n T_\sigma(f)|^2 \right)^{\frac{1}{2}} \right\|_{L^p} \\ & \leq C_p(n) K \left\| \left(\sum_{j_1, \dots, j_n \in \mathbb{Z}} \left[M^{(1)} \dots M^{(n)} (|\Delta_{j_1}^{\theta, 1} \dots \Delta_{j_n}^{\theta, n} f|^\rho) \right]^{\frac{2}{\rho}} \right)^{\frac{1}{2}} \right\|_{L^p} \\ & \leq C_p(n) K \left\| \left(\sum_{j_1, \dots, j_n \in \mathbb{Z}} |\Delta_{j_1}^{\theta, 1} \dots \Delta_{j_n}^{\theta, n} f|^2 \right)^{\frac{1}{2}} \right\|_{L^p} \\ & \leq C_p(n) K \|f\|_{L^p}. \end{aligned}$$

Notice that the second inequality follows from Lemma 2.1 and the third inequality is obtained by applying the Fefferman-Stein inequality [6] on the Lebesgue space $L^{\frac{p}{2}}$ in

each of the variables y_1, \dots, y_n . Observe that the Fefferman-Stein inequality makes use of the assumptions $2/\rho > 1$ and $p/2 > 1$.

The case $1 < p < 2$ follows by a duality argument, while the case $p = 2$ is a consequence of Plancherel's theorem and of a Sobolev embedding into L^∞ . \square

3. THE PROOF OF THEOREM 1.1; AN INTERPOLATION ARGUMENT

When $p = 2$ no derivatives are required of σ for T_σ to be bounded. To mitigate the effect of the requirement of the derivatives of σ for T_σ to be bounded on L^p for $p \neq 2$, we apply an interpolation argument between $p = 2$ and $p = 1 + \delta$.

We shall use the notation introduced at the beginning of the previous section, and we shall denote

$$\Gamma(\{s_\ell\}_{\ell=1}^n) = \Gamma(s_1, \dots, s_n) = (I - \partial_1^2)^{\frac{s_1}{2}} \cdots (I - \partial_n^2)^{\frac{s_n}{2}}.$$

The following result will be the key interpolation estimate:

Theorem 3.1. *Fix $1 < p_0, p_1, r_0, r_1 < \infty$, $0 < s_1^0, \dots, s_n^0, s_1^1, \dots, s_n^1 < \infty$. Suppose that $r_0 s_\ell^0 > 1$ and $r_1 s_\ell^1 > 1$ for all $\ell = 1, \dots, n$. Let ψ be as before. Assume that for $k \in \{0, 1\}$ we have*

$$(3.8) \quad \|T_\sigma(f)\|_{L^{p_k}} \leq K_k \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| \Gamma(s_1^k, \dots, s_n^k) \left[\sigma(2^J \xi) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^{r_k}} \|f\|_{L^{p_k}}$$

for all $f \in \mathcal{C}_0^\infty(\mathbb{R}^n)$. For $0 < \theta < 1$ and $\ell = 1, \dots, n$ define

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{r} = \frac{1-\theta}{r_0} + \frac{\theta}{r_1}, \quad s_\ell = (1-\theta)s_\ell^0 + \theta s_\ell^1.$$

Then there is a constant C_* such that for all $f \in \mathcal{C}_0^\infty(\mathbb{R}^n)$ we have

$$(3.9) \quad \|T_\sigma(f)\|_{L^p} \leq C_* K_0^{1-\theta} K_1^\theta \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| \Gamma(s_1, \dots, s_n) \left[\sigma(2^J \xi) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^r} \|f\|_{L^p}.$$

Assuming Theorem 3.1, we complete the proof of Theorem 1.1 as follows:

Proof. Given $1 \leq r < \infty$ and $\gamma_\ell > 1/r$, $\ell = 1, \dots, n$, fix $p \in (1, \infty)$ satisfying (1.4). In fact, we can assume that $p \in (1, 2)$, since the case $p \in (2, \infty)$ follows by duality and the case $p = 2$ is a consequence of Plancherel's theorem and of a Sobolev embedding into L^∞ . In addition, assume first that $\min_\ell \gamma_\ell \leq \frac{1}{2}$. In view of (1.4), there is $\tau \in (0, 1)$ such that

$$(3.10) \quad \frac{1}{p} - \frac{1}{2} < \tau \min_\ell \gamma_\ell.$$

Set $p_1 = \frac{2}{\tau+1}$, $r_1 = 2r \min_\ell \gamma_\ell$ and $\gamma_\ell^1 = \frac{1}{2} + \varepsilon$, $\ell = 1, \dots, n$, where $\varepsilon > 0$ is a real number whose exact value will be specified later. Since $p_1 > 1$ and $r_1 \gamma_\ell^1 > 2\gamma_\ell^1 > 1$, $\ell = 1, \dots, n$, Proposition 2.2 yields that

$$(3.11) \quad \|T_\sigma(f)\|_{L^{p_1}} \leq C_1 \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| \Gamma(\gamma_1^1, \dots, \gamma_n^1) \left[\sigma(2^J \xi) \prod_{\ell=1}^n \widehat{\psi}_\ell(\xi_\ell) \right] \right\|_{L^{r_1}} \|f\|_{L^{p_1}}.$$

Pick $p_0 = 2$. Let θ be the real number satisfying

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1},$$

namely, $\theta = \frac{2}{\tau}(\frac{1}{p} - \frac{1}{2})$. Observe that, by (3.10), $0 < \theta < 2 \min_\ell \gamma_\ell \leq 1$. Finally, choose real numbers r_0 and γ_ℓ^0 , $\ell = 1, \dots, n$, in such a way that

$$(3.12) \quad \frac{1}{r} = \frac{1-\theta}{r_0} + \frac{\theta}{r_1}, \quad \gamma_\ell = (1-\theta)\gamma_\ell^0 + \theta\gamma_\ell^1.$$

We claim that, for a suitable choice of $\varepsilon > 0$, one has $r_0 > 2$ and $r_0\gamma_\ell^0 > 1$, $\ell = 1, \dots, n$. Indeed, since $\min_\ell \gamma_\ell \leq \frac{1}{2}$, we have $r_1 \leq r$, and thus, by (3.12), $r_0 \geq r \geq r_1 > 2$. Further,

$$r_0\gamma_\ell^0 = \frac{r_1 r (\gamma_\ell - \theta\gamma_\ell^1)}{r_1 - \theta r} = \frac{r \min_k \gamma_k (\gamma_\ell - \frac{\theta}{2} - \theta\varepsilon)}{\min_k \gamma_k - \frac{\theta}{2}} \geq \frac{r \min_k \gamma_k (\min_k \gamma_k - \frac{\theta}{2} - \theta\varepsilon)}{\min_k \gamma_k - \frac{\theta}{2}}.$$

Since $r \min_k \gamma_k > 1$ and $\min_k \gamma_k - \frac{\theta}{2} > 0$, one gets $r_0\gamma_\ell^0 > 1$ if $\varepsilon > 0$ is small enough. Consequently, the space $\{g : \Gamma(\gamma_1^0, \dots, \gamma_n^0)g \in L^{r_0}\}$ embeds in L^∞ , and we thus have

$$(3.13) \quad \|T_\sigma(f)\|_{L^2} \leq C_1 \sup_{j_1, \dots, j_n \in \mathbb{Z}} \left\| \Gamma(\gamma_1^0, \dots, \gamma_n^0) \left[\sigma(2^J \xi) \prod_{\ell=1}^n \widehat{\psi}_\ell(\xi_\ell) \right] \right\|_{L^{r_0}} \|f\|_{L^2}.$$

The boundedness of T_σ on $L^p(\mathbb{R}^n)$ for any σ satisfying (1.3) thus follows from Theorem 3.1. Finally, if $\min_\ell \gamma_\ell > \frac{1}{2}$, then the required assertion follows directly from Proposition 2.2. \square

Proof of Theorem 3.1. The proof of Theorem 3.1 follows closely that of [1, Theorem 4.7] and for this reason we only provide an outline of its proof with few details. Throughout the proof we shall use the notation introduced at the beginning of the previous section. Also, whenever $J \in \mathbb{Z}^n$, we denote

$$\varphi_J = \Gamma(s_1, \dots, s_n) \left[\sigma(2^J \xi) \widehat{\psi}(\xi) \right],$$

and for z with real part in $[0, 1]$ we define

$$(3.14) \quad \sigma_z(\xi) = \sum_{J \in \mathbb{Z}^n} \Gamma\left(\{-s_\ell^0(1-z) - s_\ell^1 z\}_{\ell=1}^n\right) \left[|\varphi_J|^{r(\frac{1-z}{r_0} + \frac{z}{r_1})} e^{i \operatorname{Arg}(\varphi_J)} \right] (2^{-J} \xi) \widehat{\psi}(2^{-J} \xi).$$

For any given $\xi \in \mathbb{R}^n$, this sum has only finitely many terms and one can show that

$$(3.15) \quad \|\sigma_{\tau+it}\|_{L^\infty} \lesssim (1 + |t|)^{\frac{3n}{2}} \left(\sup_{J \in \mathbb{Z}^n} \|\Gamma(s_1, \dots, s_n)[\sigma(2^J \xi) \widehat{\psi}(\xi)]\|_{L^r} \right)^{\frac{r}{r_\tau}},$$

where r_τ is the real number satisfying $\frac{1}{r_\tau} = \frac{1-\tau}{r_0} + \frac{\tau}{r_1}$.

Let T_z be the family of operators associated to the multipliers σ_z . Fix $f, g \in \mathcal{C}_0^\infty$ and $1 < p_0 < p < p_1 < \infty$. Given $\varepsilon > 0$ there exist functions f_z^ε and g_z^ε such that $\|f_z^\varepsilon - f\|_{L^p} < \varepsilon$, $\|g_z^\varepsilon - g\|_{L^{p'}}$ $< \varepsilon$, and that

$$\begin{aligned} \|f_{it}^\varepsilon\|_{L^{p_0}} &\leq (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p_0}}, & \|f_{1+it}^\varepsilon\|_{L^{p_1}} &\leq (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p_1}}, \\ \|g_{it}^\varepsilon\|_{L^{p'_0}} &\leq (\|g\|_{L^{p'}}^{p'} + \varepsilon)^{\frac{1}{p'_0}}, & \|g_{1+it}^\varepsilon\|_{L^{p'_1}} &\leq (\|g\|_{L^{p'}}^{p'} + \varepsilon)^{\frac{1}{p'_1}}. \end{aligned}$$

The existence of f_z^ε and g_z^ε is folklore and is omitted; for a similar construction see [1, Theorem 3.3]. Let $F(z) = \int T_{\sigma_z}(f_z^\varepsilon)g_z^\varepsilon dx$. Then $F(z)$ is equal to

$$\begin{aligned} &\int_{\mathbb{R}^n} \sigma_z(\xi) \widehat{f}_z^\varepsilon(\xi) \widehat{g}_z^\varepsilon(\xi) d\xi \\ &= \sum_{J \in \mathbb{Z}^n} \int_{\mathbb{R}^n} \Gamma\left(\{-s_\ell^0(1-z) - s_\ell^1 z\}_{\ell=1}^n\right) \left[|\varphi_J|^{r\left(\frac{1-z}{r_0} + \frac{z}{r_1}\right)} e^{i\text{Arg}(\varphi_J)} \right] (2^{-J}\xi) \widehat{\theta}(2^{-J}\xi) \widehat{f}_z^\varepsilon(\xi) \widehat{g}_z^\varepsilon(\xi) d\xi \\ &= \sum_{J \in \mathbb{Z}^n} \int_{\mathbb{R}^n} \left[|\varphi_J|^{r\left(\frac{1-z}{r_0} + \frac{z}{r_1}\right)} e^{i\text{Arg}(\varphi_J)} \right] (2^{-J}\xi) \Gamma\left(\{-s_\ell^0(1-z) - s_\ell^1 z\}_{\ell=1}^n\right) \left[\widehat{\theta}(2^{-J}\cdot) \widehat{f}_z^\varepsilon \widehat{g}_z^\varepsilon \right](\xi) d\xi. \end{aligned}$$

The function $F(z)$ is analytic on the strip $0 < \Re(z) < 1$ and continuous up to the boundary. Notice that $\sigma_{it}(2^K \cdot) \widehat{\psi}$ picks up only the terms of (3.14) for which J differs from K in some coordinate by at most one unit. For simplicity we may therefore take $K = J$ in the calculation below. Using the Kato-Ponce inequality we may “remove” the factor $\widehat{\theta}$ and write

$$\begin{aligned} &\|T_{\sigma_{it}}(f_{it}^\varepsilon)\|_{L^{p_0}} \\ &\leq K_0 \sup_{K \in \mathbb{Z}^n} \left\| \Gamma(s_1^0, \dots, s_n^0) [\sigma_{it}(2^K \cdot) \widehat{\psi}] \right\|_{L^{r_0}} \|f_{it}^\varepsilon\|_{L^{p_0}} \\ &\leq K_0 \sup_{K \in \mathbb{Z}^n} \left\| \Gamma(\{s_\ell^0 - s_\ell^0(1-it) - s_\ell^1 it\}_{\ell=1}^n) [|\varphi_K|^{r\left(\frac{1-it}{r_0} + \frac{it}{r_1}\right)} e^{i\text{Arg}(\varphi_K)}] \right\|_{L^{r_0}} \|f_{it}^\varepsilon\|_{L^{p_0}} \\ &\lesssim (1 + |t|)^{\frac{3n}{2}} K_0 \sup_{K \in \mathbb{Z}^n} \|\varphi_K\|_{L^r}^{\frac{r}{r_0}} (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p_0}}. \end{aligned}$$

Using Hölder’s inequality $|F(it)| \leq \|T_{\sigma_{it}}(f_{it}^\varepsilon)\|_{L^{p_0}} \|g_{it}^\varepsilon\|_{L^{p'_0}}$, we may therefore write

$$|F(it)| \leq C(1 + |t|)^{\frac{3n}{2}} K_0 \sup_{J \in \mathbb{Z}^n} \left\| \Gamma(\{s_\ell^0\}_{\ell=1}^n) [\sigma(2^J \cdot) \widehat{\psi}] \right\|_{L^r}^{\frac{r}{r_0}} (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p_0}} (\|g\|_{L^{p'}}^{p'} + \varepsilon)^{\frac{1}{p'_0}}$$

for some constant $C = C(n, r_0, s_\ell^0, s_\ell^1)$. Similarly, for some constant $C = C(n, r_1, s_\ell^0, s_\ell^1)$ we obtain

$$|F(1 + it)| \leq C(1 + |t|)^{\frac{3n}{2}} K_1 \sup_{J \in \mathbb{Z}^n} \|\Gamma(\{s_\ell\}_{\ell=1}^n)[\sigma(2^J \cdot) \widehat{\psi}]\|_{L^r}^{\frac{r}{r_1}} (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p_1}} (\|g\|_{L^{p'}}^{p'} + \varepsilon)^{\frac{1}{p_1}}.$$

Thus for $z = \tau + it$, $t \in \mathbb{R}$ and $0 \leq \tau \leq 1$ it follows from (3.15) and from the definition of $F(z)$ that

$$|F(z)| \leq C''(1 + |t|)^{\frac{3n}{2}} \left(\sup_{J \in \mathbb{Z}^n} \|\Gamma(s_1, \dots, s_n)[\sigma(2^J \cdot) \widehat{\psi}]\|_{L^r} \right)^{\frac{r}{r_1}} \|f_z^\varepsilon\|_{L^2} \|g_z^\varepsilon\|_{L^2} = A_\tau(t),$$

noting that $\|f_z^\varepsilon\|_{L^2} \|g_z^\varepsilon\|_{L^2}$ is bounded above by constants independent of t and τ . Since $A_\tau(t) \leq \exp(Ae^{a|t|})$, the hypotheses of three lines lemma are valid. It follows that

$$|F(\theta)| \leq C K_0^{1-\theta} K_1^\theta \sup_{J \in \mathbb{Z}^n} \|\Gamma(\{s_\ell\}_{\ell=1}^n)[\sigma(2^J \cdot) \widehat{\psi}]\|_{L^r} (\|f\|_{L^p}^p + \varepsilon)^{\frac{1}{p}} (\|g\|_{L^{p'}}^{p'} + \varepsilon)^{\frac{1}{p}}.$$

Taking the supremum over all functions $g \in L^{p'}$ with $\|g\|_{L^{p'}} \leq 1$, a simple density argument yields for some $C_* = C_*(n, r_1, r_2, s_\ell^0, s_\ell^1)$

$$\|T_\sigma(f)\|_{L^p} \leq C_* K_0^{1-\theta} K_1^\theta \sup_{J \in \mathbb{Z}^n} \|\Gamma(s_1, \dots, s_n)[\sigma(2^J \cdot) \widehat{\psi}]\|_{L^r} \|f\|_{L^p}.$$

This completes the proof of the sufficiency part of Theorem 1.1. The proof of the necessity part is postponed to section 5. \square

4. THE PROOF OF THEOREM 1.2

Crucial ingredients needed for the proof of Theorem 1.2 are two one-dimensional inequalities contained in the following lemma.

Lemma 4.1. *Let ψ be as in Theorem 1.2. If $k \in \mathbb{Z}$, $\gamma > 0$ and $1 < r < \infty$ are such that $\gamma r > 1$, then*

$$(4.16) \quad \|f(2^k \cdot) \widehat{\psi}\|_{L^r(\mathbb{R})} \leq C \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})}$$

and

$$(4.17) \quad \|(-\partial^2)^{\frac{\gamma}{2}} [f(2^k \cdot) \widehat{\psi}]\|_{L^r(\mathbb{R})} \leq C(1 + 2^{k(\gamma - \frac{1}{r})}) \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})}.$$

Proof. Since $\gamma r > 1$, the Sobolev embedding theorem yields

$$(4.18) \quad |f(2^k x)| \leq \|f\|_{L^\infty(\mathbb{R})} \leq C \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})} \quad \text{for a.e. } x \in \mathbb{R}.$$

Therefore,

$$\|f(2^k \cdot) \widehat{\psi}\|_{L^r(\mathbb{R})} \leq C \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})} \|\widehat{\psi}\|_{L^r(\mathbb{R})} = C' \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})}.$$

This proves (4.16).

Further, using the Kato-Ponce inequality [10], the estimate (4.18) and the fact that $\widehat{\psi}$ is smooth and with compact support, we obtain

$$\begin{aligned}
& \|(-\partial^2)^{\frac{\gamma}{2}} [f(2^k \cdot) \widehat{\psi}]\|_{L^r(\mathbb{R})} \\
& \leq C \left(\|(-\partial^2)^{\frac{\gamma}{2}} [f(2^k \cdot)]\|_{L^r(\mathbb{R})} \|\widehat{\psi}\|_{L^\infty(\mathbb{R})} + \|f(2^k \cdot)\|_{L^\infty(\mathbb{R})} \|(-\partial^2)^{\frac{\gamma}{2}} \widehat{\psi}\|_{L^r(\mathbb{R})} \right) \\
& \leq C \left(\|(-\partial^2)^{\frac{\gamma}{2}} [f(2^k \cdot)]\|_{L^r(\mathbb{R})} + \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})} \right) \\
& = C \left(2^{k(\gamma - \frac{1}{r})} \|(-\partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})} + \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})} \right) \\
& \leq C (2^{k(\gamma - \frac{1}{r})} + 1) \|(I - \partial^2)^{\frac{\gamma}{2}} f\|_{L^r(\mathbb{R})},
\end{aligned}$$

namely, (4.17). \square

Proof of Theorem 1.2. Set $F(\xi) = \sum_{a=-n}^n \widehat{\Phi}(2^a \xi)$, $\xi \in \mathbb{R}^n$. Then $F(\xi) = 1$ for any ξ satisfying $\frac{1}{2^n} \leq |\xi| \leq 2^n$. Therefore, if j_1, \dots, j_n are integers and $j := \max\{j_1, \dots, j_n\}$, then $F(2^{j_1-j} \xi_1, \dots, 2^{j_n-j} \xi_n) = 1$ on $\{(\xi_1, \dots, \xi_n) : \frac{1}{2} \leq |\xi_1| \leq 2, \dots, \frac{1}{2} \leq |\xi_n| \leq 2\}$. Consequently,

$$\prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) = F(2^{j_1-j} \xi_1, \dots, 2^{j_n-j} \xi_n) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell).$$

Using this, we can write

$$\begin{aligned}
& \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^r} \\
& = \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n) F(2^{j_1-j} \xi_1, \dots, 2^{j_n-j} \xi_n) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^r} \\
& \leq C \sum_{a=-n}^n \sum_{\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}} \\
& \left\| (-\partial_{i_1}^2)^{\frac{\gamma_{i_1}}{2}} \cdots (-\partial_{i_k}^2)^{\frac{\gamma_{i_k}}{2}} \left[\sigma(2^{j_1} \xi_1, \dots, 2^{j_n} \xi_n) \widehat{\Phi}(2^{j_1-j+a} \xi_1, \dots, 2^{j_n-j+a} \xi_n) \prod_{\ell=1}^n \widehat{\psi}(\xi_\ell) \right] \right\|_{L^r}.
\end{aligned}$$

Using the estimate (4.17) in variables i_1, \dots, i_k and inequality (4.16) in the remaining variables, we estimate the corresponding term in the last expression by a constant multiple of

$$\begin{aligned}
& \left[\prod_{s=1}^n (1 + 2^{(j_{i_s} - j + a)(\gamma_{i_s} - \frac{1}{r})}) \right] \left\| (I - \partial_1^2)^{\frac{\gamma_1}{2}} \cdots (I - \partial_n^2)^{\frac{\gamma_n}{2}} \left[\sigma(2^{j-a} \xi) \widehat{\Phi}(\xi) \right] \right\|_{L^r} \\
& \leq C (1 + 2^{n \max_{\ell=1, \dots, n} (\gamma_\ell - \frac{1}{r})})^n \left\| (I - \Delta)^{\frac{\gamma_1 + \dots + \gamma_n}{2}} \left[\sigma(2^{j-a} \xi) \widehat{\Phi}(\xi) \right] \right\|_{L^r}
\end{aligned}$$

$$\leq C \sup_{m \in \mathbb{Z}} \left\| (I - \Delta)^{\frac{\gamma_1 + \dots + \gamma_n}{2}} \left[\sigma(2^m \xi) \widehat{\Phi}(\xi) \right] \right\|_{L^r}.$$

This implies (1.5). \square

5. EXAMPLES AND REMARKS

Next we discuss examples that indicate the sharpness of Theorem 1.1. As stated, the sufficient condition presented in Theorem 1.1 is optimal in the sense that if T_σ is bounded from $L^p(\mathbb{R}^n)$ to itself for all σ satisfying (1.3), then (1.4) holds. This has been observed, at least in the two-dimensional case with both smoothness parameters equal, by Carbery and Seeger [3, remark after Proposition 6.1]. We provide an example in the spirit of theirs, given by an explicit closed-form expression and valid in all dimensions $n \geq 2$.

Example 5.1. *Given $\alpha \in (0, 1)$, consider the function*

$$\sigma(\xi, \eta) = \varphi(|\eta|) e^{-\frac{\xi^2}{2}} |\eta|^{i\xi} (\log |\eta|)^{-\alpha}, \quad (\xi, \eta) \in \mathbb{R} \times \mathbb{R}^{n-1} = \mathbb{R}^n,$$

where φ is a smooth function on the line such that $0 \leq \varphi \leq 1$, $\varphi = 0$ on $(-\infty, 8]$ and $\varphi = 1$ on $[9, \infty)$. Then

- (i) σ satisfies (1.3), with r large enough, whenever $\gamma_1 = \alpha$ and $\gamma_2, \dots, \gamma_n$ are arbitrary positive real numbers;
- (ii) σ is an L^p Fourier multiplier for a given $1 < p < \infty$ if and only if $\alpha > |\frac{1}{p} - \frac{1}{2}|$.

The previous example indicates that condition (1.3) does not guarantee that T_σ is L^p bounded unless all indices $\gamma_1, \dots, \gamma_n$ in (1.3) are larger than $|\frac{1}{p} - \frac{1}{2}|$. In particular, for a given $i \in \{1, \dots, n\}$, one does not have boundedness on the critical line $\gamma_i = |\frac{1}{p} - \frac{1}{2}|$, no matter how large the remaining parameters are.

Let us now verify the statement of part (i) of Example 5.1. We shall first prove that

$$(5.19) \quad \sup_{k, \ell \in \mathbb{Z}} \|(I - \partial_\xi^2)^{\frac{s}{2}} (I - \Delta_\eta)^{\frac{s}{2}} [\widehat{\psi}(\xi) \widehat{\Phi}(\eta) \sigma(2^k \xi, 2^\ell \eta)]\|_{L^r} < \infty$$

for any $s > 0$ and $r > 1$. Here, Φ denotes a Schwartz function on \mathbb{R}^{n-1} whose Fourier transform is supported in the set $\{\eta \in \mathbb{R}^{n-1} : \frac{1}{2} \leq |\eta| \leq 2\}$ and which satisfies $\sum_{\ell \in \mathbb{Z}} \widehat{\Phi}(2^\ell \eta) = 1$ for all $\eta \neq 0$. Indeed, for any $k, \ell \in \mathbb{Z}$, $\ell \geq 3$, and for any given nonnegative integer m , we have

$$\|\widehat{\psi}(\xi) \widehat{\Phi}(\eta) \sigma(2^k \xi, 2^\ell \eta)\|_{L^r} \leq C \ell^{-\alpha}$$

and

$$\|(I - \partial_\xi^2)^{\frac{1}{2}} (I - \Delta_\eta)^{\frac{m}{2}} [\widehat{\psi}(\xi) \widehat{\Phi}(\eta) \sigma(2^k \xi, 2^\ell \eta)]\|_{L^r} \leq C \ell \cdot \ell^{-\alpha},$$

where the constant C is independent of k and ℓ . Interpolating between these two estimates, we obtain

$$\|(I - \partial_\xi)^{\frac{\alpha}{2}}(I - \Delta_\eta)^{\frac{\alpha m}{2}}[\widehat{\psi}(\xi)\widehat{\Phi}(\eta)\sigma(2^k\xi, 2^\ell\eta)]\|_{L^r} \leq C.$$

Notice also that the last inequality in fact holds for all integers k, ℓ , since the function $\widehat{\psi}(\xi)\widehat{\Phi}(\eta)\sigma(2^k\xi, 2^\ell\eta)$ is identically equal to 0 if $\ell \leq 2$. Hence, we have

$$\sup_{k, \ell \in \mathbb{Z}} \|(I - \partial_\xi)^{\frac{\alpha}{2}}(I - \Delta_\eta)^{\frac{\alpha m}{2}}[\widehat{\psi}(\xi)\widehat{\Phi}(\eta)\sigma(2^k\xi, 2^\ell\eta)]\|_{L^r} \leq C,$$

and interpolating between variants of this estimate corresponding to different values of m , we obtain (5.19) for any $s > 0$. Now, part (i) of Example 5.1 follows by an application of Theorem 1.2 in the variable η .

Let us finally focus on part (ii) of Example 5.1. If $\alpha > |\frac{1}{p} - \frac{1}{2}|$, then σ is an L^p Fourier multiplier thanks to (i) and Theorem 1.1. Let us now prove that T_σ is not L^p bounded if $\alpha \leq |\frac{1}{p} - \frac{1}{2}|$. By duality, it suffices to discuss only the case when $1 < p < 2$. Further, by a result of Herz and Rivière [12], our claim will follow if we show that T_σ is not bounded on the mixed norm space $L^p(\mathbb{R}; L^2(\mathbb{R}^{n-1}))$. (We take first the L^2 norm in η and then the L^p norm in ξ .)

Let f be the function on \mathbb{R}^n whose Fourier transform satisfies

$$\widehat{f}(\xi, \eta) = e^{-\frac{\xi^2}{2}} \varphi(|\eta|) |\eta|^{\frac{1-n}{2}} (\log |\eta|)^{-\frac{1}{2}} (\log \log |\eta|)^{-\beta}, \quad (\xi, \eta) \in \mathbb{R} \times \mathbb{R}^{n-1}.$$

Using the Plancherel theorem in the variable η , it is easy to check that $f \in L^p(\mathbb{R}; L^2(\mathbb{R}^{n-1}))$ whenever $\beta > \frac{1}{2}$. Our next goal is to prove that $T_\sigma f = \mathcal{F}^{-1}(\sigma \widehat{f})$ does not belong to $L^p(\mathbb{R}; L^2(\mathbb{R}^{n-1}))$ if $\beta \in (\frac{1}{2}, \frac{1}{p}]$. Using Plancherel's theorem in the variable η again, this is equivalent to showing that $\mathcal{F}_\xi^{-1}(\sigma \widehat{f})$ is not in $L^p(\mathbb{R}; L^2(\mathbb{R}^{n-1}))$, where \mathcal{F}_ξ^{-1} stands for the inverse Fourier transform in the ξ variable.

Observe that

$$\begin{aligned} \mathcal{F}_\xi^{-1}(\sigma \widehat{f})(x, \eta) &= C e^{-\frac{1}{4}(2\pi x + \log |\eta|)^2} \varphi^2(|\eta|) |\eta|^{\frac{1-n}{2}} (\log |\eta|)^{-\alpha - \frac{1}{2}} (\log \log |\eta|)^{-\beta} \\ &\geq C \chi_{\{(x, \eta): x < -2, e^{-2\pi x - 1} < |\eta| < e^{-2\pi x}\}}(x, \eta) e^{2\pi x \frac{n-1}{2}} (-x)^{-\alpha - \frac{1}{2}} (\log(-x))^{-\beta}. \end{aligned}$$

Therefore,

$$\|\mathcal{F}_\xi^{-1}(\sigma \widehat{f})\|_{L^p(\mathbb{R}; L^2(\mathbb{R}^{n-1}))} \geq C \left(\int_{-\infty}^{-2} (-x)^{(-\alpha - \frac{1}{2})p} (\log(-x))^{-\beta p} dx \right)^{\frac{1}{p}} = \infty,$$

which yields the desired conclusion.

Acknowledgment: We would like to thank Andreas Seeger for his useful comments.

REFERENCES

- [1] A. P. Calderón, A. Torchinsky, *Parabolic maximal functions associated with a distribution, II*, Adv. in Math. **24** (1977), 101–171.
- [2] A. Carbery, *Differentiation in lacunary directions and an extension of the Marcinkiewicz multiplier theorem*, Ann. de l'Institut Fourier **38** (1988), 157–168.
- [3] A. Carbery and A. Seeger, *H^p - and L^p -Variants of multiparameter Calderón-Zygmund theory*, Trans. AMS **334** (1992), 719–747.
- [4] A. Carbery and A. Seeger, *Homogeneous Fourier multipliers of Marcinkiewicz type*, Ark. Mat. **33** (1995), 45–80.
- [5] J. Duoandikoetxea, *Fourier Analysis*, Grad. Studies in Math., Vol 29, AMS, Providence, RI, 2001.
- [6] C. Fefferman, E. M. Stein, *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107–115.
- [7] L. Grafakos, *Classical Fourier Analysis*, 3rd edition, GTM 249, Springer-Verlag, NY 2014.
- [8] L. Grafakos, D. He, P. Honzík, H. V. Nguyen, *The Hörmander multiplier theorem I: The linear case revisited*, Illinois J. Math. **61** (2017), 25–35.
- [9] L. Grafakos, D. He, H. V. Nguyen, L. Yan, *Multilinear multiplier theorems and applications*, J. Fourier. Anal. Appl., to appear.
- [10] L. Grafakos, S. Oh, *The Kato-Ponce inequality*, Comm. in PDE **39** (2014), 1128–1157.
- [11] L. Grafakos, L. Slavíková, *A sharp version of the Hörmander multiplier theorem*, Int. Math. Res. Not. IMRN, to appear.
- [12] C. Herz, N. M. Rivière, *Estimates for translation invariant operators on spaces with mixed norms*, Studia Math. **44** (1972), 511–515.
- [13] I. I. Jr. Hirschman, *On multiplier transformations*, Duke Math. J. **26** (1959), 221–242.
- [14] A. Miyachi, *On some Fourier multipliers for $H^p(\mathbb{R}^n)$* , J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), 157–179.
- [15] A. Miyachi, N. Tomita, *Minimal smoothness conditions for bilinear Fourier multipliers*, Rev. Mat. Iberoam. **29** (2013), 495–530.
- [16] L. Slavíková. *On the failure of the Hörmander multiplier theorem in a limiting case*, Rev. Mat. Iberoam., to appear.
- [17] S. Wainger, *Special trigonometric series in k -dimensions*, Mem. Amer. Math. Soc. **59** (1965), 1–102.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA
E-mail address: grafakosl@missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA
E-mail address: slavikoval@missouri.edu