

Pseudodifferential operators with homogeneous symbols

Loukas Grafakos*
Department of Mathematics
University of Missouri
Columbia, MO 65211

Rodolfo H. Torres†
Department of Mathematics
University of Kansas
Lawrence, KS 66045

Abstract

We prove boundedness of pseudodifferential operators with symbols satisfying the conditions $|\partial_\xi^\beta \partial_x^\gamma a(x, \xi)| \leq C_{\beta, \gamma} |\xi|^{m-|\beta|+|\gamma|}$ on homogeneous Besov-Lipschitz and Triebel-Lizorkin spaces

1 Introduction

The study of pseudodifferential operators with symbols in the exotic classes $S_{1,1}^m$ has received a lot of attention. These are operators of the form

$$(Tf)(x) = \int_{\mathbf{R}^n} e^{ix \cdot \xi} a(x, \xi) \hat{f}(\xi) d\xi,$$

where the symbol $a(x, \xi)$ is a $C^\infty(\mathbf{R}^n \times \mathbf{R}^n)$ function satisfying

$$|\partial_\xi^\beta \partial_x^\gamma a(x, \xi)| \leq C_{\beta, \gamma} (1 + |\xi|)^{m-|\beta|+|\gamma|},$$

for all β, γ n -tuples of nonnegative integers. The interest in such operators is in part due to the role they play in the paradifferential calculus of Bony

⁰*AMS subject classification numbers:* Primary 42B20, 47G30. Secondary 46E35, 35B65.

⁰*Key words and phrases:* Pseudodifferential operators, homogeneous symbols, Besov-Lipschitz spaces, Triebel-Lizorkin spaces

*Supported in part by NSF grant DMS 9623120 and the University of Missouri Research Board.

†Supported in part by NSF grants DMS 9623251 and DMS 9696267.

[1]. The fact that not all such operators of order zero are bounded on L^2 complicates their study. Nevertheless, the exotic pseudodifferential operators do preserve spaces of smooth functions. See for example Meyer [12], Runst [15], Bourdaud [2], as well as Stein [16] and the references therein.

The continuity results are often obtained by making use of the so-called singular integral realization of the operators. This involves proving estimates on the Schwartz kernels of the pseudodifferential operators similar to those of the kernels of Calderón–Zygmund operators. There is, however, an alternative approach working directly with the symbols of the pseudodifferential operators. This approach has been pursued by Hörmander in [9] and [10] for L^2 -based Sobolev spaces. The ideas in those papers combined with wavelets techniques were latter extended to L^p -based Sobolev spaces and other more general spaces of smooth functions by Torres [17].

In this note we consider C^∞ symbols $a(x, \xi)$ in $\mathbf{R}^n \times (\mathbf{R}^n \setminus \{\mathbf{0}\})$ that satisfy the following conditions: For all n -tuples of nonnegative integers β and γ there exist positive constants $C_{\beta, \gamma}$ such that

$$|\partial_\xi^\beta \partial_x^\gamma a(x, \xi)| \leq C_{\beta, \gamma} |\xi|^{m-|\beta|+|\gamma|}, \quad (1)$$

for $(x, \xi) \in \mathbf{R}^n \times (\mathbf{R}^n \setminus \{\mathbf{0}\})$. We call such symbols homogeneous symbols of type $(1, 1)$ and order m . The class of all such symbols will be denoted by $\dot{S}_{1,1}^m$.

Our purpose is to show boundedness for pseudodifferential operators with symbols in $\dot{S}_{1,1}^m$ on homogeneous function spaces. Our results are motivated by a theorem of Grafakos [8] who proved boundedness of pseudodifferential operators with symbols in $\dot{S}_{1,1}^m$ on homogeneous Lipschitz spaces. The proof in [8] follows more or less the singular integral approach of Stein in [16]. In this paper we use a wavelet approach borrowing ideas from [17].

The appropriate setting for our results is in the context of the homogeneous Triebel-Lizorkin $\dot{F}_p^{\alpha, q}(\mathbf{R}^n)$ spaces and Besov-Lipschitz spaces $\dot{B}_p^{\alpha, q}(\mathbf{R}^n)$ (see the definition of these spaces below). As with inhomogeneous symbols, it is possible to show that the Schwartz kernels of the operators of order zero satisfy for $x \neq y$ estimates of the form

$$|\partial_x^\gamma K(x, y)| + |\partial_y^\gamma K(x, y)| \leq C_\gamma |x - y|^{-n-|\gamma|} \quad (2)$$

and even better estimates for $|x - y|$ large. See for example the book by Meyer [13], p.294. It is then possible to analyze boundedness properties of the operators using versions of the T1-Theorem of David and Journé [4]. Moreover, such type of results are applied to pseudodifferential operators

with inhomogeneous symbols in the book of Meyer mentioned before (p.329) in the context of Besov spaces with smoothness $\alpha > 0$ and $p, q > 1$ (cf. also [11]). The arguments therein may be adapted to pseudodifferential operators with homogeneous symbols and the same range of parameters for the Besov spaces. Our approach, however, will be based on some very simple calculations that notoriously work for both full scales of $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ and $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ spaces.

Let $\mathcal{S}(\mathbf{R}^n)$ be the space of Schwartz test functions and denote its dual by $\mathcal{S}'(\mathbf{R}^n)$, the space of tempered distributions. In this paper the Fourier transform of a function $f \in \mathcal{S}(\mathbf{R}^n)$ is given by $\hat{f}(\xi) = \int f(x)e^{-ix \cdot \xi} dx$ and $\mathcal{S}_0(\mathbf{R}^n)$ is used to denote the subspace of $\mathcal{S}(\mathbf{R}^n)$ consisting of all functions whose Fourier transform vanishes to infinite order at zero. The dual space of $\mathcal{S}_0(\mathbf{R}^n)$, with respect to the topology inherited from $\mathcal{S}(\mathbf{R}^n)$, is $\mathcal{S}'/\mathcal{P}(\mathbf{R}^n)$, the set of tempered distributions modulo polynomials. The Triebel-Lizorkin and Besov-Lipschitz spaces are defined as follows. Let φ be a function in $\mathcal{S}(\mathbf{R}^n)$ satisfying $\text{supp } \hat{\varphi} \subset \{\xi : 1/2 \leq |\xi| \leq 2\}$ and $|\hat{\varphi}| \geq C > 0$ for $3/5 \leq |\xi| \leq 5/3$. Define $\varphi_j(\xi) = 2^{jn}\varphi(2^j\xi)$. For α real, $0 < p, q < \infty$, and f in $\mathcal{S}'/\mathcal{P}(\mathbf{R}^n)$ define the Triebel-Lizorkin and Besov-Lipschitz norms of f by

$$\|f\|_{\dot{F}_p^{\alpha,q}(\mathbf{R}^n)} = \|(\sum_{j=-\infty}^{\infty} (2^{j\alpha}|\varphi_j * f|)^q)^{1/q}\|_{L^p},$$

and

$$\|f\|_{\dot{B}_p^{\alpha,q}(\mathbf{R}^n)} = (\sum_{j=-\infty}^{\infty} (\|2^{j\alpha}(\varphi_j * f)\|_{L^p})^q)^{1/q},$$

respectively. Note the the above definitions are given modulo all polynomials, so strictly speaking an element of the spaces $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ and $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ is an equivalent class of distributions. It can be shown that the definition of these spaces is independent of φ . The space $\mathcal{S}_0(\mathbf{R}^n)$ is dense in all of these spaces. For these and further properties of these function spaces we refer to the books [14] and [19].

We have the following results for operators with symbols in $\dot{S}_{1,1}^m$ acting on the spaces $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ and $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$.

Theorem 1.1 *Let $0 < p, q < \infty$. For $\alpha > n(\max(1, p^{-1}, q^{-1}) - 1)$, every pseudodifferential operator*

$$(Tf)(x) = \int_{\mathbf{R}^n} a(x, \xi) \hat{f}(\xi) e^{ix \cdot \xi} d\xi, \quad f \in \mathcal{S}_0$$

with symbol $a(x, \xi)$ in $\dot{S}_{1,1}^m$ extends to a bounded operator that maps the space $\dot{F}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$. For $\alpha \leq n(\max(1, p^{-1}, q^{-1}) - 1)$, every T as above which satisfies $T^*x^\gamma = 0$ for all $|\gamma| \leq n(\max(1, p^{-1}, q^{-1}) - 1) - \alpha$ extends to a bounded operator from $\dot{F}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$.

Theorem 1.2 *Let $0 < p, q < \infty$. For $\alpha > n(\max(1, p^{-1}, q^{-1}) - 1)$, every pseudodifferential operator*

$$(Tf)(x) = \int_{\mathbf{R}^n} a(x, \xi) \hat{f}(\xi) e^{ix \cdot \xi} d\xi, \quad f \in \mathcal{S}_0 \quad (3)$$

with symbol $a(x, \xi)$ in $\dot{S}_{1,1}^m$ extends to a bounded operator that maps the space $\dot{B}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$. For $\alpha \leq n(\max(1, p^{-1}, q^{-1}) - 1)$, every T as above which satisfies $T^*x^\gamma = 0$ for all $|\gamma| \leq n(\max(1, p^{-1}, q^{-1}) - 1) - \alpha$ extends to a bounded operator from $\dot{B}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$.

We end this section with the following observation. Note that if $m - |\gamma| + |\beta| < 0$, then $\partial_\xi^\gamma \partial_x^\beta a(x, \xi)$ is singular at $\xi = 0$ and for a general function f in \mathcal{S} the integral in (3) is not absolutely convergent. For this reason it is natural to define the operator T initially on \mathcal{S}_0 .

2 The proof of the theorems

As discussed above it is natural to consider T initially defined on \mathcal{S}_0 . Moreover, we have

Lemma 2.1 *Let $a(x, \xi)$ be a symbol in $\dot{S}_{1,1}^m$. Then the pseudodifferential operator with symbol $a(x, \xi)$ maps \mathcal{S}_0 to \mathcal{S} . In particular, its formal transpose T^* maps \mathcal{S}' to \mathcal{S}'/\mathcal{P} .*

Let f be a function in \mathcal{S}_0 and let Δ_ξ be the Laplace operator in the variable ξ . Since for any positive integer N ,

$$(I - \Delta_\xi)^N (e^{ix \cdot \xi}) = (1 + |x|^2)^N e^{ix \cdot \xi},$$

an integration by parts gives

$$(Tf)(x) = \int_{\mathbf{R}^n} e^{ix \cdot \xi} \frac{(I - \Delta_\xi)^N}{(1 + |x|^2)^N} (a(x, \xi) \hat{f}(\xi)) d\xi. \quad (4)$$

Since \hat{f} vanishes to infinity order at the origin, the conditions on the symbol $a(x, \xi)$ and an application of Leibnitz's rule give

$$|(Tf)(x)| \leq \frac{\mathcal{R}_{m,N}(f)}{(1 + |x|^2)^N}, \quad (5)$$

where $\mathcal{R}_{m,N}$ is an appropriate seminorm in \mathcal{S} . A similar computation applies to the derivatives $\partial^\gamma(Tf)$, thus proving the lemma.

For an n -tuple of integers k and an integer j denote by Q_{jk} the dyadic cube $\{(x_1, \dots, x_n) \in \mathbf{R}^n : k_i \leq 2^j x_i < k_i + 1\}$, by $x_{Q_{jk}}$ its "lower left corner" $2^{-j}k$ and by $l(Q_{jk})$ its side length 2^{-j} . For Q dyadic let $\varphi_Q(x) = |Q|^{1/2} \varphi_j(x - x_Q)$. A function φ as in the definition of the homogeneous spaces can be chosen so that for $f \in \mathcal{S}'$,

$$f = \sum_Q \langle f, \varphi_Q \rangle \varphi_Q, \quad (6)$$

where $\langle f, \varphi_Q \rangle$ simply denotes the action of the distribution f on the test function φ_Q . For f in $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ or $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ the convergence in (6) is in the (quasi)-norm of the spaces and for f in \mathcal{S}_0 is in the topology of \mathcal{S} . See [6] and [7].

It follows from Lemma 2.1 that the action of a pseudodifferential operator on \mathcal{S}_0 can be expressed as

$$Tf = \sum_Q \langle f, \varphi_Q \rangle T\varphi_Q. \quad (7)$$

The operator given by (7) is the one that is extended to the whole homogeneous space in our Theorems. We now turn to the proofs.

The map

$$S_\varphi(f) = \{\langle f, \varphi_Q \rangle\}_Q \quad (8)$$

is called the φ -transform (or sequence of nonorthogonal wavelet coefficients). It is well known by the work in [5], [6] that the homogeneous φ -transform provides a characterization of the spaces $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ and $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ via the equivalence of norms

$$\|f\|_{\dot{F}_p^{\alpha,q}(\mathbf{R}^n)} \sim \left\| \left(\sum_j \left(\sum_{l(Q)=2^{-j}} |Q|^{-\alpha/n-1/2} |\langle f, \varphi_Q \rangle| \chi_Q \right)^q \right)^{1/q} \right\|_{L^p}$$

and

$$\|f\|_{\dot{B}_p^{\alpha,q}(\mathbf{R}^n)} \sim \left(\sum_j \left\| \sum_{l(Q)=2^{-j}} |Q|^{-\alpha/n-1/2} \langle f, \varphi_Q \rangle |\chi_Q| \right\|_{L^p}^q \right)^{1/q},$$

where χ_Q is the characteristic function of Q .

For α , p , and q as above let $J = n / \min(1, p, q)$ and let $[\alpha]$ be the integer part in α . A smooth molecule for $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ or $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ associated with a dyadic cube Q with side length $l(Q)$ is a function m_Q satisfying:

$$\int x^\gamma m_Q(x) dx = 0 \quad \text{if } |\gamma| \leq [J - n - \alpha] \quad (9)$$

$$|m_Q(x)| \leq |Q|^{-1/2} (1 + l(Q)^{-1} |x - x_Q|)^{-\max(J+\varepsilon, J+\varepsilon-\alpha)} \quad (10)$$

$$|\partial^\gamma m_Q(x)| \leq |Q|^{-1/2-|\gamma|/n} (1 + l(Q)^{-1} |x - x_Q|)^{-J-\varepsilon}, \quad |\gamma| \leq [\alpha] + 1 \quad (11)$$

The importance of these functions is due to the fact that if

$$f = \sum_Q s_Q m_Q$$

in \mathcal{S}' , where $\{m_Q\}$ is a family of smooth molecules for $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ or $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$, then

$$\|f\|_{\dot{F}_p^{\alpha,q}(\mathbf{R}^n)} \leq C \left\| \left(\sum_j \left(\sum_{l(Q)=2^{-j}} |Q|^{-\alpha/n-1/2} |s_Q| |\chi_Q| \right)^q \right)^{1/q} \right\|_{L^p}$$

or

$$\|f\|_{\dot{B}_p^{\alpha,q}(\mathbf{R}^n)} \leq C \left(\sum_j \left\| \sum_{l(Q)=2^{-j}} |Q|^{-\alpha/n-1/2} |s_Q| |\chi_Q| \right\|_{L^p}^q \right)^{1/q}.$$

For the above results we refer again to [6] and [7].

Now, let T be a linear continuous operator from $\mathcal{S}_0 \rightarrow \mathcal{S}'$. Assume that

$$T\varphi_Q = C|Q|^{-m/n} m_Q, \quad (12)$$

where $\{m_Q\}_Q$ is a family of smooth molecules for $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ or $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$. Then using the φ -transform it is easy to see that T can be extended as a bounded operator from $\dot{F}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ or from $\dot{B}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$.

Suppose that T is a pseudodifferential operator whose symbol $a(x, \xi)$ is in $\dot{S}_{1,1}^m$. By the remark (12) above it will suffice to show that for a fixed dyadic cube Q , $T\varphi_Q$ is a scaled multiple of a molecule. A simple dilation shows that

$$(T\varphi_Q)(x) = \int e^{ix \cdot \xi} a(x, \xi) \hat{\varphi}_Q(\xi) d\xi = 2^{jn/2} (T_Q \varphi)(2^j x - k), \quad (13)$$

if $Q = Q_{jk}$, where we set

$$(T_Q f)(x) = \int e^{ix \cdot \xi} a(2^{-j}(x+k), 2^j \xi) \hat{f}(\xi) d\xi.$$

Let us fix a multi-index γ . We have

$$(\partial^\gamma T_Q \varphi)(x) = \int_{\mathbf{R}^n} e^{ix \cdot \xi} \sum_{\delta \leq \gamma} C_\delta (i\xi)^\delta \partial_x^{\gamma-\delta} (a(2^{-j}(x+k), 2^j \xi) \hat{\varphi}(\xi)) d\xi \quad (14)$$

for certain C_δ constants, where $\delta \leq \gamma$ simply means that $\delta_j \leq \gamma_j$ for all $j = 1, \dots, n$.

Now fix $N > \max(J, J - \alpha)/2$. An integration by parts gives

$$(\partial^\gamma T_Q \varphi)(x) = \int_{\mathbf{R}^n} e^{ix \cdot \xi} \frac{(I - \Delta_\xi)^N}{(1 + |x|^2)^N} \sum_{\delta \leq \gamma} C_\delta (i\xi)^\delta \partial_x^{\gamma-\delta} (a(2^{-j}(x+k), 2^j \xi) \hat{\varphi}(\xi)) d\xi \quad (15)$$

By Leibnitz's rule, there exist constants $K_{\alpha,\beta}$ such that

$$\begin{aligned} (I - \Delta_\xi)^N \partial_x^{\gamma-\delta} (a(2^{-j}(x+k), 2^j \xi) \hat{\varphi}(\xi) (i\xi)^\delta) = \\ \sum_{|\alpha+\beta|=2N} K_{\alpha,\beta} \partial_\xi^\beta \partial_x^{\gamma-\delta} (a(2^{-j}(x+k), 2^j \xi) \partial_\xi^\alpha (\hat{\varphi}(\xi) (i\xi)^\delta)). \end{aligned}$$

Using the estimates (1) we conclude that

$$\begin{aligned} |\partial_\xi^\beta \partial_x^{\gamma-\delta} (a(2^{-j}(x+k), 2^j \xi))| &\leq C_{\beta,\gamma-\delta} 2^{j|\beta|} 2^{-j(|\gamma|-|\delta|)} |2^j \xi|^{m+(|\gamma|-|\delta|)-|\beta|} \\ &\leq C 2^{jm} |\xi|^{m+(|\gamma|-|\delta|)-|\beta|}. \end{aligned}$$

Summing over all α , β , and δ as above and using that $|\xi| \sim 1$ we conclude that the integrand in (15) is pointwise bounded by $C_\gamma 2^{jm} (1 + |x|^2)^{-N}$ and thus

$$|(\partial^\gamma T_Q \varphi)(x)| \leq C_\gamma 2^{jm} (1 + |x|^2)^{-N}. \quad (16)$$

We now dilate and translate (16) to deduce that

$$\begin{aligned} |(\partial^\gamma T_Q \varphi)(x)| &\leq C 2^{jn/2} 2^{j|\gamma|} 2^{jm} (1 + |2^j x - k|^2)^{-N} \\ &\leq C |Q|^{-m/n-1/2-|\gamma|/n} (1 + l(Q)^{-1} |x - x_Q|)^{-\max(J+\varepsilon, J+\varepsilon-\alpha)} \end{aligned}$$

We must now check the vanishing moment condition for $T\varphi_Q$. If $[J - n - \alpha] < 0$ this condition is vacuous. If $[J - n - \alpha] \geq 0$, this is an easy consequence of the hypothesis $T^* x^\gamma = 0$, since

$$\int x^\gamma (T\varphi_Q)(x) dx = \langle x^\gamma, T\varphi_Q \rangle = \langle T^*(x^\gamma), \varphi_Q \rangle = 0.$$

Both theorems are now proved.

We conclude this section with some remarks.

1. For $p > 1$ let p' denote $p/(p-1)$. It can be shown that the spaces $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$ can actually be considered as spaces of distributions modulo polynomials of degree less than or equal to $[\alpha - n/p]$. (See [6] page 154.) For $1 < p, q < \infty$, if T maps $\dot{F}_p^{\alpha+m,q}(\mathbf{R}^n)$ to $\dot{F}_p^{\alpha,q}(\mathbf{R}^n)$, then by duality T^* maps $\dot{F}_{p'}^{-\alpha,q'}(\mathbf{R}^n)$ to $\dot{F}_{p'}^{-\alpha-m,q'}(\mathbf{R}^n)$. It follows that for T^* to be even well-defined on $\dot{F}_{p'}^{-\alpha,q'}(\mathbf{R}^n)$ it must annihilate polynomials of degree $[-\alpha - n/p']$. The conditions on T^* in the second part of Theorems 1.1 and 1.2 for $\alpha < 0$ become then necessary for $p \rightarrow 1^+$.

2. For more general operators T with kernels satisfying estimates (2) and the usual weak boundedness property assumed in the T1-Theorem (which is always satisfied by pseudodifferential operators of order zero and their transposes), the results in [13] state that the conditions $T(x^\gamma) = 0$ for all $|\gamma| \leq [\alpha]$ imply that T is bounded on $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ for $\alpha > 0$ and $p, q \geq 1$. Let now $\alpha < 0$. If T is a pseudodifferential operator in $\dot{S}_{1,1}^0$, then T^* is not necessarily a pseudodifferential operator. Nevertheless, its kernel $K^*(x, y)$ is $K(y, x)$ and, hence, it still satisfies the estimates (2) by symmetry. The results in [13] state that if T^* annihilates polynomials of degree less than or equal to $[-\alpha]$, then T^* is bounded on $\dot{B}_p^{-\alpha,q}$, and then by duality $T = T^{**}$ is bounded in $\dot{B}_p^{\alpha,q}$, which agrees with our results. Such duality arguments are not available for other values of the parameters but our proof is still valid.

3. When $m \neq 0$, we could have precomposed the operator with an appropriate power of the Laplacian and reduced our proof to the case $m = 0$. There are also versions of the T1-Theorem for general operators whose kernels satisfy appropriate m -versions of the estimates (2). In principle, such results could be applied to operators in $\dot{S}_{1,1}^m$ and some values of the

parameters α , p and q , but they lead to weaker results than the ones we have presented here (cf. [18]).

4. Note that our approach does not require the giving of a precise meaning on the action of an operator T (satisfying (2)) on polynomials, as it is usually required in the T1-Theorem and its versions. We only need to know that T^* acts on polynomials, which is automatic by duality.

5. Operators with homogeneous symbols of degree m in ξ satisfying estimates (1) were studied by Calderón and Zygmund [3]. For this subclass of symbols a partial calculus holds but it does not extend to the whole class $\dot{S}_{1,1}^m$ (see [16], p.268).

3 Examples and applications

Symbols in $\dot{S}_{1,1}^m$ which are independent of x exist in abundance. For instance, it is easy to see that the reciprocal of an elliptic polynomial of n variables which is homogeneous of degree $m > 0$ is in $\dot{S}_{1,1}^{-m}$.

An example of a homogeneous symbol in the class $\dot{S}_{1,1}^0$ is the following:

$$\sum_{k=-\infty}^{+\infty} e^{i2^k x \cdot \xi} \phi(2^{-k} \xi),$$

where ϕ is a smooth bump supported away from the origin. More generally, suppose that the sequence of smooth functions $\{m_k(x)\}_{k \in \mathbf{Z}}$ in \mathbf{R}^n satisfies

$$\|\partial^\beta m_k\|_\infty \leq C_\alpha 2^{|\alpha|k}, \quad (17)$$

for all α n -tuples of nonnegative integers and k any integer. Then the symbol

$$a(x, \xi) = \sum_{k=-\infty}^{+\infty} m_k(x) \phi(2^{-k} \xi)$$

is in $\dot{S}_{1,1}^0$.

We now give an application. Let Δ_j be the Littlewood-Paley operators defined by $\widehat{\Delta_j g}(\xi) = \widehat{g}(\xi) \phi(2^{-j} \xi)$, where ϕ is a smooth bump supported away from the origin which satisfies $\sum_{j \in \mathbf{Z}} \phi(2^{-j} \xi) = 1$ for all $\xi \neq 0$.

Suppose now that f is a function on \mathbf{R}^n that satisfies

$$\sum_{j \in \mathbf{Z}} \|\Delta_j f\|_\infty \leq C(f) < \infty. \quad (18)$$

Let F be a C^∞ function on \mathbf{R}^n with $F(0) = 0$. Suppose that f is in some of the homogeneous function space discussed in the previous section with index of smoothness $\alpha > 0$. Denote such space by $X_p^{\alpha,q}$. We claim that $F(f)$ lies in the same space $X_p^{\alpha,q}$. For the proof of this we borrow the ideas of Bony as presented in [12].

For k integer define

$$f_k = \sum_{j=-\infty}^k \Delta_j f, \quad (19)$$

and write $f = \lim_{k \rightarrow \infty} f_k$, with uniform convergence because of (18). The functions f_k have Fourier transforms with compact support and they are smooth (actually analytic of exponential type). Moreover, by (18), they are uniformly bounded. Using (18) and Bernstein's inequality we obtain the following estimates for their derivatives

$$\|\partial^\alpha f_k\|_\infty \leq C_\alpha 2^{|\alpha|k} \|f_k\|_\infty \leq C(f) C_\alpha 2^{|\alpha|k}. \quad (20)$$

Write now

$$F(f) = \lim_{N \rightarrow \infty} \sum_{k=-N}^N F(f_k) - F(f_{k-1}) = \sum_{k=-\infty}^{+\infty} F(f_k) - F(f_{k-1}), \quad (21)$$

where convergence is justified from the fact that F is continuous, $F(0) = 0$ and that $f_N \rightarrow f$ and $f_{-N} \rightarrow 0$ uniformly as $N \rightarrow +\infty$. Next apply the mean value theorem to write (21) as

$$F(f)(x) = \sum_{k=-\infty}^{+\infty} m_k(x) (\Delta_k f)(x),$$

where

$$m_k = \int_0^1 F'(tf_k + (1-t)f_{k-1}) dt. \quad (22)$$

Using (20), the smoothness of F , and the chain rule, we see that the functions m_k satisfy (17). We conclude that the symbol

$$a(x, \xi) = \sum_{j=-\infty}^{+\infty} m_j(x) \phi(2^{-j}\xi)$$

is in $\dot{S}_{1,1}^0$. We have that $F(f) = T_a f$. It follows from Theorems 1.1 and 1.2 that the function $F(f)$ is in $X_p^{\alpha,q}$, provided $\alpha > 0$ and $p, q \geq 1$, or if

$\alpha > n(\max(1, p^{-1}, q^{-1}) - 1)$. Observe that because of the nonlinearity of the problem, in the estimate

$$\|F(f)\|_{X_p^{\alpha,q}} \leq C_f \|f\|_{X_p^{\alpha,q}}, \quad (23)$$

the constant C_f depends on f . In fact, both the functions m_k and the symbol a depend on f . If we assume that $F(t) = t^D$ then, after a careful examination of the arguments above and of the proof of the theorems in the previous section, we see that C_f in (23) is controlled by a suitable (large) power of the bound $C(f)$ in (18).

Finally observe that the left hand side of (18) is the $\dot{B}_\infty^{0,1}$ norm of f . By some well-known facts about functions of exponential type (see [19]),

$$\|\Delta_j f\|_\infty \leq C 2^{jn/p} \|\Delta_j f\|_p.$$

From this one obtains the inequality (Sobolev-Besov embedding)

$$\sum_{j \in \mathbf{Z}} \|\Delta_j f\|_\infty \leq C \sum_{j \in \mathbf{Z}} 2^{jn/p} \|\Delta_j f\|_p.$$

Then, in particular, the application described can be used in $\dot{B}_p^{\alpha,q}(\mathbf{R}^n)$ with $\alpha = n/p$ and $q = 1$ yielding (23) with C_f controlled by a power of $\|f\|_{\dot{B}_p^{n/p,1}}$.

References

- [1] J. Bony, *Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires*, Ann. Sci. Ecole Norm. Sup. 14 (1981), 209–256.
- [2] G. Bourdaud, *Une algèbre maximale d'opérateurs pseudodifférentiels*, Comm. Partial Diff. Eq. 13 (1988), 1059–1083.
- [3] A. Calderón and A. Zygmund, *Singular integral operators and differential equations*, Amer. J. Math. 79 (1957), 801–821.
- [4] G. David and J-L. Journé *A boundedness criterion for generalized Calderón–Zygmund operators*, Ann. of Math. 120 (1984), 371–397.
- [5] M. Frazier, B. Jawerth, *Decomposition of Besov spaces*, Indiana Univ. Math. J. 34 (1985), 777–799.

- [6] M. Frazier, B. Jawerth, *A discrete transform and decompositions of distribution spaces*, J. Func. Anal. 93(1) (1990), 34–169.
- [7] M. Frazier, B. Jawerth, and G. Weiss, *Littlewood-Paley theory and the study of function spaces*, CBMS Regional Conference Series in Mathematics 79, 1991.
- [8] L. Grafakos, *Boundedness of pseudodifferential operators on homogeneous Lipschitz spaces*, submitted.
- [9] L. Hörmander, *Pseudodifferential operators of type 1,1*, Comm. Partial Diff. Eq. 13 (1988) 1085–1111.
- [10] L. Hörmander, *Continuity of pseudodifferential operators of type 1,1*, Comm. Partial Diff. Eq. 14 (1989) 231–243.
- [11] P. Lemarié, *Continuité sur les espaces de Besov des opérateurs définis par des intégrales singulières*, Ann. Inst. Fourier 35 (1985), 175–187.
- [12] Y. Meyer, *Regularité des solutions des équations aux dérivées partielles non linéaires*, Springer-Verlag Lectures Notes in Math. 842 (1980), 293–302
- [13] Y. Meyer, *Ondelettes et opérateurs II*, Hermann Ed., Paris, France, 1990.
- [14] J. Peetre, *New thoughts on Besov spaces*, Duke University Math. Series 1, Durham, 1976.
- [15] T. Runst, *Pseudo-differential operators in Hardy-Triebel spaces*, Z. Anal. Anw. 2 (1983), 235–242.
- [16] E. Stein, *Harmonic Analysis, Real Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, Princeton, New Jersey, 1993.
- [17] R. Torres, *Continuity properties of pseudodifferential operators of type (1,1)*, Comm. Par. Diff. Eq. 15 (9) (1990), 1313–1328.
- [18] R. Torres, *Boundedness results for operators with singular kernels on distribution spaces*, Mem. Amer. Math. Soc., 442 (1991).
- [19] H. Triebel, *Theory of function spaces*, Monographs in Mathematics, Vol. 78, Birkhauser Verlag, Basel, 1983.